

The effect of the COVID-19 pandemic on professional and advanced cycling athletes: a cross-sectional study in Poland

Pawel Rzymiski^{1,2*}, Piotr Rzymiski^{3,4}, Bartłomiej Burzynski⁵, Tomasz Jurys⁵, Michalina Knapik⁶, Izabela Rzymaska⁷

¹Department of Mothers' and Child's Health, Poznan University of Medical Sciences, Gynecologic and Obstetrical University Hospital, Poznan, Poland

²University Center for Sport and Medical Studies, Poznan University of Medical Sciences, Poznan, Poland

³Department of Environmental Medicine, Poznan University of Medical Sciences, Poznan, Poland

⁴Integrated Science Association (ISA), Universal Scientific Education and Research Network (USERN), Poznan, Poland

⁵Department of Rehabilitation, Faculty of Health Sciences, Medical University of Silesia, Katowice, Poland

⁶Doctoral School, Department of Urology, Faculty of Medical Sciences in Zabrze, Medical University of Silesia in Katowice, Poland

⁷Department of Social and Humanistic Science, Poznan University of Medical Sciences, Poznan, Poland

***Corresponding author:**

Pawel Rzymiski
Department of Mothers' and Child's Health
Poznan University of Medical Sciences,
Gynecologic and Obstetrical University Hospital
33 Polna St
60-535 Poznan, Poland
E-mail: pawel@rzymiski.eu

Submitted: 8 August 2022; **Accepted:** 7 October 2022

Online publication: 28 October 2022

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/155211>

Copyright © 2022 Termedia & Banach

Abstract

Introduction: The COVID-19 pandemic has affected different communities, but data on its impact on cyclists are scarce.

Material and methods: The present study assessed the training patterns of professional and advanced amateur cycling athletes in Poland ($n = 130$) during the COVID-19 pandemic compared with the pre-pandemic season and in relation to their epidemiological situation and quality of life factors.

Results: The volume and weekly frequency of training were not decreased during the pandemic. However, their high-intensity interval training time (weekly 31.6 vs. 37.0 min, $p = 0.004$) and maximum weekly Training Stress Score (660 vs. 715, $p = 0.001$) decreased. Compared to uninfected cyclists, those who contracted SARS-CoV-2 registered a higher score in the somatic, social and environmental domains of the WHOQOL questionnaire (56.1 vs. 50.6, $p = 0.024$; 74.1 vs. 63.6, $p = 0.032$ and 72.5 vs. 65.9, $p = 0.018$, respectively); no changes were observed in the psychologic domain. Change in the weekly frequency of training, volume, time spent in > 90% and 80–90% of HRmax (min) correlated positively only with the somatic domain of the WHOQOL. The training pattern of cyclists who experienced the death of a relative due to COVID-19 remained unaffected, but they presented a lower psychological domain in the WHOQOL (64.5 vs. 69.2, $p = 0.02$).

Conclusions: Professional and advanced amateur cyclists maintained their volume and frequency of training but reduced the intensity during the pandemic compared to the 2019 season. Changes were more likely to affect their physical status than their psychogenic condition. The pandemic had a limited effect on the training pattern of cycling athletes.

Key words: COVID-19, cycling athletes, training patterns, high-intensity interval training, SARS-CoV-2.

Introduction

The human coronavirus disease 2019 (COVID-19) led to a global lockdown and increased public anxiety and fears [1, 2]. As estimated, at least 4 billion people were forced to quarantine themselves. Observational studies indicate that such isolation has caused psychological distress, anxiety, insomnia, and depressive symptoms [3, 4]. Moreover, numerous investigations have reported adverse changes to the diet in a relatively high number of isolated individuals [5, 6]. According to some studies, the quarantine reinforced pre-existing alterations in body weight and adversely affected physical activity [5, 7]. The obligatory use of face masks also induces higher perioral humidity, temperature and breathing difficulties, which could be another limiting factor for athletes [8]. The potential infection among athletes could also influence the team training habits and the change in disinfection and protection protocols. Especially indoor athletes should be aware of air contamination with viral particles and use air disinfection similarly to the dental clinics [9]. An increase in alcohol consumption, particularly in individuals experiencing higher stress, was reported [5, 10, 11]. These findings highlight that the COVID-19 pandemic has directly affected lifestyle behaviors, raising concerns from both an individual and public health perspective.

However, the above-mentioned effects of the pandemic may not necessarily apply to those practicing sports on a regular basis. To help them cope with the epidemiological situation, some strategies encompassing personalized conditioning training have been developed [12]. At the same time, data on how the pandemic has affected the physical activities of groups practicing particular sports are scarce [13]. Cycling athletes may be particularly affected as outdoor activities have been temporarily sanctioned due to lockdowns.

In athletes under one session of heavy training, both innate and acquired immunity are decreased transiently during the recovery period after prolonged heavy exertion (≥ 90 min), typically by 15–70% [14, 15]. Empirical evidence, albeit in a small number of athletes, indicates that international athletes suffer fewer, not greater, upper respiratory infection episodes compared to national-level athletes [16, 17]. The manipulation of training volume and/or intensity includes: gradually increasing volume and intensity by 5–10% per week; increasing the frequency of shorter, spike training sessions rather than enduring fewer but longer sessions, recoveries after intensive sessions and easier weeks after 3-week blocks [18]. Psychological stress has a well-known and marked influence on immunity; thus, it plays a role in the decrease in immunity with prolonged

heavy exercise and heavy training [14]. Athletes may experience psychological stress associated with competition (or lack of competition), injury, travel, and sleep disruption; in addition, they may also experience psychological stress in their personal life related to relationship difficulties, financial hardship and bereavement. Unfortunately, psychological stress has rarely been reported in such studies; therefore, it is difficult to assess the strength of such a relationship. Some supporting evidence is, however, emerging, as more recent research shows that the level of anxiety and perceived stress (life stress in the last month) reported before exercise can significantly modify the immune response to subsequent exercise [18]. Sports practice in elite athletes should protect against psychological disorders, as stated by the International Olympic Committee (IOC), and its antidepressant effect probably disappears after ceasing to exercise [19].

Infectious diseases are usually successfully managed among athletes, but COVID-19 brought new and unprecedented challenges to the sports community, including cyclists [14, 20]. How did the epidemiological COVID-19 situation and risk of severe infection affect the training pattern, volume, and intensity? Was it influenced by psychological factors, including an athlete's becoming infected with SARS-CoV-2 or the infection of related individuals? We hypothesized that athletes could change their training patterns and the pandemic negatively influences their quality of life with the resolution of SARS-Cov-2 infection. So far, these aspects have not been subjected to any reported research.

This study aimed to provide the first answers to the above questions by assessing the training activities and patterns (frequency, volume, intensity), quality of life factors, and the epidemiological situation among professional and advanced amateur cycling athletes during the COVID-19 pandemic in Poland.

Material and methods

To explore the effect of the COVID-19 pandemic on the training activities of professional and advanced cyclists in Poland, an anonymous online survey was performed. We based our study on adult males and females from the Polish Cycling Association elite and amateur license list ($n = 1241$). Given the size of the target population, it was calculated using Cochran's formula [21] that 134 eligible individuals should be surveyed to reach the margin level of 8% at the confidence interval of 95%. Such a research approach has been recommended to swiftly reach a specific group of individuals, ensuring their safety under pandemic conditions [22]. The study was performed be-

tween 16 and 30 November 2020 and approved by the Bioethics Committee of Poznan University of Medical Sciences (KB – 170/21). Strict legal regulations limited the cycling season of 2020 under the pandemic in Poland. The first case of COVID-19 in Poland was identified on 4 March 2020; from 24 March, a nationwide lockdown was imposed; on 20 April, the ban on the recreational use of forests and parks was lifted; on 4 May, hotels and shopping centers were permitted to reopen. No cycling activities were possible between 24 March and 20 April, since cycling was only possible with the face covered. Nevertheless, all the cycling events, including professional and amateur competitions, were canceled until 31 July.

The online questionnaire (Google Forms) employed in this study was advertised and linked among Facebook professional and amateur cycling groups. A total of 40 target groups were searched on Facebook social media and were listed in two advertising posts distributed by our scientific and sports cooperators.

The inclusion criteria included: active cyclists who perform structured or unstructured training programs monitored by heart rate monitors. When available, additional data were requested: VO_2 max expressed as mL of oxygen uptake per kg of body weight per minute (minimum difference: 1.0) and Training Stress Score estimated by appropriate software (iE TrainingPeaks, Golden Cheetah and others, minimum difference: 1.0). Cyclists were recruited from advanced amateurs who do not compete, amateurs who attend races and professional cyclists. Because of different training measuring devices and software as well as different training conceptions and planning, the study focused on heart rate monitors and average derivative information.

The employed questionnaire aimed to assess:

- Training load during the COVID-19 pandemic in comparison to the pre-pandemic period (2019 season): weekly frequency and volume of sessions, weekly time spent in high-intensity intervals (expressed in minutes above 90% of HRmax, all times minimum difference 1 min), time spent in intensive intervals (expressed in minutes between 80 and 90 of HRmax), Training Stress Score (a composite number that takes into account the duration and intensity of a workout [23]) and VO_2 max expressed as ml of oxygen uptake per kg of body weight per min (when available for 2020, $n = 51$).
- SARS-CoV-2 infection history in the studied cycling athletes, death of relative due to COVID-19, and the potential effect of these circumstances on training parameters.
- Quality of life using the WHOQOL-BREF questionnaire in the following domains: physical

health, psychological, environmental, and social relationships in association with training patterns and epidemiological situations of cycling athletes.

The demographic data on each surveyed individual included gender, age, body mass index (BMI; calculated from reported weight and height), years of training (weight, height, years minimum difference 1 kg, 1 cm, 1 year), type of cycling (road, mountain bike, triathlon, etc.).

The respondents completed the WHOQOL-BREF 26-item questionnaire in the valid Polish version [24–26], an abbreviated version of WHOQOL-100 developed by the quality of life researchers at the World Health Organization. It enables an assessment of the quality of life profile in the physical health, psychological, social relationships and environment domains. The domain scores were transformed into a linear scale between 0 and 100 following the scoring guidelines [27]. A higher score indicated a better QOL. The psychometric properties of the original version are considered as very good (Cronbach's $\alpha = 0.89$).

Data from Google Forms were exported to a spreadsheet file. The training characteristics data were analyzed simultaneously for the 2019 and 2020 seasons. Changes in training frequency, volume, intensity and Training Stress Score (TSS) were calculated by subtracting the 2020 value from the 2019 value. The TSS was calculated according to the following formula: $TSS = (t \times NP \times IF) / (FTP \times 3600) \times 100$, where t is the duration of the exercise bout, NP is its normalized power, FTP is the individual's functional threshold power, IF is the intensity factor (the ratio between the NP and the individual's FTP) [24].

Statistical analysis

The statistical analysis was performed with SigmaPlot 11.0.0.77 (Systat Soft, USA). All data are presented as means \pm SD and median values or percentages. Since the variables expressed on the interval scale did not meet the assumption of a Gaussian distribution, non-parametric statistical tests were applied. The differences in studied parameters between subgroups were analyzed with the Mann-Whitey U test. Correlations were assessed with Spearman's correlation coefficient (R_s). A value of $p < 0.05$ was considered statistically significant.

Results

The online questionnaires were filled in by 131 respondents. One form was missing basic training data and was withdrawn from further analysis. The demographic breakdown of the studied groups is given in Table I.

Table I. Demographic breakdown of the studied group of cycling athletes in Poland ($n = 130$)

Parameter	<i>N</i>	%
Male/female	117/13	90/10
Age [years]	34.4 ±11.7	-
Advanced amateur (not competing)	31	23.9
Advanced amateur (competing)	74	56.9
Professional cyclist	25	19.2
Type of cycling (multiple choice possible)		
Road	106	81.5
Time trial (TT)	21	16.2
Mountain bike (MTB)	80	61.5
Enduro	1	0.8
Triathlete	21	16.2
Gravel	24	18.5
SARS-CoV-2 negative during 2020	108	83.1
SARS-CoV-2 positive during 2020	22	16.9
Subjective evaluation:		
No influence on training capacity	6	27.3%
Decreased training capacity	8	36.4%
Decreased training capacity and recovered to previous capacity	6	27.3%
Increased training capacity	2	9.0%
Do you know a person who died because of COVID-19?		
Yes	83	63.8%
No	47	36.2%
	Mean ± SD	
Age	34.4 ±11.7	
WHOQOL-BREF somatic domain (transformed)	51.1 ±11.2	
WHOQOL-BREF psychological domain (transformed)	66.2 ±12.2	
WHOQOL-BREF social domain (transformed)	65.5 ±20.7	
WHOQOL-BREF environmental domain (transformed)	67.0 ±12.7	

Training patterns

During the 2020 cycling season, 16.9% ($n = 22$) of athletes were confirmed to be SARS-CoV-2 positive, whereas 63.8% ($n = 83$) stated that they knew someone who had died due to COVID-19. The characteristics of training patterns in 2019 and 2020 are presented in Table II. Compared to the pre-pandemic season, no changes in weight, BMI, frequency of training, and exercise volume were reported. The time spent on high-intensity interval training (HIIT, > 90% of HRmax) was reduced from 37.0 to 31.6 min/week. Furthermore, although not significant, a decrease was observed in other intensive training between 80% and 90% of HRmax. For 47 athletes, the Training Stress Score was available. The weekly mean of TSS did not differ between pre-pandemic and pandemic seasons, but the maximum weekly TSS was reduced in the latter (660.6 vs. 767.0, $p < 0.05$). This represents the reduction of higher intervals and maximum weekly TSS while retaining overall training load.

We analyzed the potential influence of a positive COVID-19 test on the training patterns and quality of life measured by WHOQOL-BREF. Subjectively, a few respondents reported a decrease in training capacity, which in eight individuals did not recover (Table I). All the SARS-CoV-2-infected athletes were asymptomatic or had mild symptoms of COVID-19. Their training patterns after the infection period did not change. A tendency to reduce training volume and intensive sessions was observed, but the data variation was likely influenced by the smaller sample size bias. Interestingly, cyclists infected with SARS-CoV-2 had better physical health, social relations domain, and environmental domain scoring, but not in the psychological domain (Table III). Athletes infected with SARS-CoV-2 seemed slightly less experienced (4.8 years vs. 7.9) and had a lower VO_2 max (54.6 vs. 60.1) than those who did not acquire the infection, but the difference was not significant. Infected cyclists reported the resolution of symptoms without severe complications.

Table II. Comparison between the pre-pandemic year 2019 and pandemic 2020 season regarding training patterns in the studied group of cycling athletes in Poland ($n = 130$)

Parameter		2019 season (prior to COVID-19)	2020 season (during COVID-19)	P-value
VO ₂ max [ml/kg/min]*	Mean	–	58.9 ±10.6	–
Years of training [years]	Mean	–	7.4 ±8.6	–
Weight [kg]	Mean	74.4 ±13.0	74.5 ±13.2	0.37
	Median	74	74	
BMI [kg/m ²]	Mean	23.2 ±3.7	23.2 ±3.8	0.32
	Median	23.1	23.2	
Frequency of training [/week]	Mean	4.4 ±1.6	4.3 ±1.9	0.73
	Median	4	4	
Training volume [h/week]	Mean	8.1 ±3.	8.0 ±4.3	0.68
	Median	8	8	
Time spent in > 90% of HRmax [min/week]	Mean	37.0 ±38.3	31.6 ±41.7	0.04
	Median	27.5	20.0	
Time spent in 80–90% of HRmax [min/week]	Mean	93.3 ±96.9	84.6 ±91.7	0.20
	Median	60.0	60.0	
TSS [average/week]*	Mean	396.8 ±216.8	391.0 ±207.8	0.15
	Median	400.0	400.0	
TSS [maximum/week]*	Mean	767.0 ±439.6	660.6 ±319.2	0.001
	Median	715.0	660.0	

*Data available for 47/130 cyclists.

Influence on quality of life

We hypothesized that infection with SARS-CoV-2 or death among family and friends due to COVID-19 could affect the training patterns. As observed, the routine training plan and the training frequency, volume, and intensity did not change statistically significantly in cyclists who experienced the death of their relatives/friends due to COVID-19 (Table IV). At the same time, such individuals revealed lower psychological health domain scoring (statistically significant) and social relation domain scoring (nearly significant) measured by WHOQOL-BREF.

Physical health correlated positively with the change of training frequency, volume and intensity. At the same time, psychological health positively affected only the frequency of training but not its volume and intensity. Age, training experience, BMI and VO₂max had no influence on the whole health report (Table V).

Discussion

To the best of our knowledge, this is the first study to report on the effects of the COVID-19 pandemic on the training patterns of cycling athletes. The epidemiological situation has caused global disruptions to the world of sports with cancellations of major and smaller-scale sporting events, including cycling competitions. By itself,

this could affect the training volume and intensity of amateur and professional cyclists.

In general, there are different strategies to achieve optimal results in cycling. Planning the whole year season is crucial for achieving the best results for priority racings (A) independent of cycling types: road, cross country, or triathlon. Several measurement parameters and devices have been developed in sport science for proper coaching or self-coaching. They are based on the conception of training frequency, volume (duration), and intensity, whereby intensity could be considered as full session intensity and one session intensity, measured at least by heart rate monitors or a power meter. However, diet, recovery, and micro/macrocycle planning are needed to achieve the highest possible efficiency. This includes low-intensity basic training mixed with an interval session, periodization or inverted periodization, polarization, high altitude training/living, and a racing calendar.

The efficacy of day-to-day high-intensity interval training (HIIT) monitored by power on a cycloergometer, initially at a minimum of 91% HRmax, was effective. Although the increase in VO₂max in all three types of training was similar to that reported in the literature, no differences or additional advantages of any of the intensive regimes were observed [28]. As demonstrated, this regime in designed percentages and intensities spent in

Table III. Changes in training and health domains among SARS-CoV-2-positive and SARS-CoV-2-negative cycling athletes in Poland ($n = 130$)

Parameter		Infected cyclists ($n = 22$)	Non-infected cyclists ($n = 108$)	P-value
Training experience [years]		4.8 \pm 4.0	7.9 \pm 9.2	0.320
BMI		23.5 \pm 2.5	23.1 \pm 3.9	0.659
VO ₂ max [ml/kg/min]*		54.6 \pm 6.9	60.1 \pm 11.2	0.28
Age [years]		35.0 \pm 9.7	34.3 \pm 12.1	0.675
Change in frequency of training [/week]	Mean	-0.18 \pm 1.39	0.03 \pm 1.73	
	Median	0	0	0.789
Change in training volume [h/week]	Mean	-0.29 \pm 3.78	0.0 \pm 3.81	
	Median	0	0	0.564
Change in time spent in > 90% of HRmax [min]	Mean	-13.9 \pm 32.5	-3.5 \pm 39.8	
	Median	-1.5	0	0.406
Change in time spent in 80–90% of HRmax [min]	Mean	-18.0 \pm 76.2	-6.5 \pm 68.5	
	Median	0	0	0.604
Change in TSS [average/week]*	Mean	-34.2 \pm 178.9	5.1 \pm 202.1	
	Median	-36	0	0.120
Change in TSS [maximum/week]*	Mean	-24.8 \pm 390.4	-102.6 \pm 406.9	
	Median	-125	-10	0.540
WHOQOL-BREF somatic domain (transformed)	Mean	56.1 \pm 12.4	50.6 \pm 10.7	
	Median	56	53	0.024
WHOQOL-BREF psychological domain (transformed)	Mean	66.0 \pm 13.0	66.3 \pm 12.1	
	Median	69	69	0.836
WHOQOL-BREF social domain (transformed)	Mean	74.1 \pm 15.1	63.6 \pm 21.2	
	Median	75	69	0.032
WHOQOL-BREF environmental domain (transformed)	Mean	72.5 \pm 11.2	65.9 \pm 12.8	
	Median	75	69	0.018

*Data available for 47/130 cyclists.

certain zones has a significant training effect on power output and VO₂max [29]. Both sprint interval training (SIT 30-s all-out bouts, together 6 min in sprint power range weekly) and high-intensity interval training (HIIT > 90% of HRmax, together 48 min/week) three times weekly are other ways to improve cycling performance. Participants improved their VO₂max by 4.5 and 4.4, respectively, and the SIT group benefited additionally from a higher sum of skinfold reduction (36.3 vs. 20.5 mm) [30].

As evaluated in the present study, all intensive or high-intensity cycling regimes, irrespective of the exact interval design and session plan, could be a cause of apprehension during the COVID-19 pandemic. Based on the observations of the present study, the professional and amateur cyclists did not significantly reduce the frequency and volume of training sessions. A decrease in HIIT was noted and a weak tendency to reduce intensive training volume (80–90% HRmax). But the overall weekly TSS did not change, suggesting that intensity was substituted partially by an increase

in low- to middle-intensity volume training. This is rational because of infection avoidance and risks described in the literature after intensive sessions [14, 15]. Roche *et al.* even reported an increase in the duration of training in 31% of athletes, with a mean increase from 92 to 103 min per day during the pandemic. But the mean heart rate remained unchanged. This partially corresponds to our results, but the results can be highly location-specific. Additionally, the cited study was performed on a very inhomogeneous group of 131 runners, cyclists and triathletes [31]. In another study conducted among 65 runners, the training pattern during the pandemic changed towards an increase in volume but the intensity did not change [32]. Muriel *et al.* reported a decrease in weekly training volumes by 25.8–52.2% in zones Z1 to Z6 except for the aerobic Z2. This caused the reduction of 5 and 20 min performances among 18 professional cyclists [33]. In contrast, elite athletes in Italy reduced their energy expenditure (MET) and hours of training. However, the studied athletes were very heterogeneous; individual elite athletes

Table IV. Changes in training patterns in the studied group of Polish cyclists ($n = 130$) in relation to whether they experienced death of family members or friends due to COVID-19

Parameter		Cyclists who knew somebody who died ($n = 83$)	Cyclists who did not know somebody who died ($n = 47$)	P-value
Change in frequency of training [/week]	Mean	0.08 ±1.8	-0.14 ±1.30	
	Median	0	0	0.57
Change in training volume [h/week]	Mean	0.00 ±4.1	-0.15 ±3.21	
	Median	0	0	0.88
Change in time spent in > 90% of HRmax [min]	Mean	-3.0 ±39.9	-9.4 ±37.2	
	Median	0	0	0.97
Change in time spent in 80–90% of HRmax [min]	Mean	-7.9 ±77.3	-9.5 ±55.4	
	Median	0	0	0.90
Change in TSS [average/week]*	Mean	-6.2 ±200.1	6.1 ±192.1	
	Median	0	-20	0.67
Change in TSS [maximum/week]*	Mean	-112.1 ±425.4	-9.2 ±318.0	
	Median	-32.5	0	0.22
WHOQOL-BREF somatic domain (transformed)	Mean	52.1 ±11.1	50.0 ±11.3	
	Median	56	50	0.28
WHOQOL-BREF psychological domain (transformed)	Mean	64.5 ±12.6	69.2 ±11.0	
	Median	69	69	0.02
WHOQOL-BREF social domain (transformed)	Mean	63.6 ±20.7	68.6 ±20.6	
	Median	69	75	0.11
WHOQOL-BREF environmental domain (transformed)	Mean	67.2 ±12.8	66.5 ±12.8	
	Median	69	69	0.62

*Data available for 47/130 cyclists.

completed a greater number of training sessions during COVID-19 compared to team athletes [19]. Both aerobic and resistance interval training is adequate for improving stable coronary heart disease [30]. Although maintaining at least low to middle intensive training could not be beneficial for cycling efficiency progress, it may better stimulate the immune system against infectious respiratory diseases. During a pandemic, reasonable training in accordance with better life hygiene (diet, sleep) should be recommended [16, 17].

Previous viral diseases, including the Middle East respiratory syndrome (MERS), have been associated with myocardial injury and myocarditis with troponin elevation, thought to be due to increased cardiac physiologic stress, hypoxia, or direct myocardial injury [34, 35]. Athletes should be aware of additionally generating hypoxia because acute heart failure can be the primary presenting manifestation of COVID-19 infection. One study found that COVID-19-induced cardiomyopathy occurs in 33% of patients with the infection [36]. Another study found that heart failure was present in 24% of patients and was associated with an increased risk of mortality, even when nearly half had no known history of hypertension or cardiovascular disease [37]. Up to 20–30% of patients hospitalized with COVID-19 have evidence

of myocardial involvement, but this percentage in subclinical or mild symptoms of disease remains unknown [38]. Athletes are a special subgroup of patients, and given the rapidly evolving knowledge base about the virus and changing governmental and public health recommendations, it is difficult to provide precise answers to this highly relevant question. For athletes suffering from flu-like symptoms, myocarditis has always been a potential complication of viral syndromes. While serum biomarkers of myocardial damage are commonly elevated in the acute stages, it is important to note that raised troponins do not equate to a diagnosis of myocarditis and may be secondary to intensive physical training. For those that have fully recovered without ongoing cardiovascular symptoms (symptom-free at least for 7 days and no sooner than day 10 from the onset of symptoms), it is recommended to carry out cardiac testing, including 12-lead ECG and echocardiography, and, if the results are abnormal or the symptoms are persistent or prolonged, additional MRI. Measuring serum cardiac troponin should be confined to athletes with ongoing symptoms compatible with myocarditis and those with imaging studies that reveal impaired myocardial function [39]. However, in rare cases, patients returning to high-level sport following confirmed myocarditis

Table V. Spearman's correlations between changes in training parameters in 2020 and WHOQOL domain scores among studied cyclists ($n = 130$)

Parameter	WHOQOL physical health domain (transformed)	WHOQOL psychological health domain (transformed)	WHOQOL social relations domain (transformed)	WHOQOL environment domain (transformed)
Change in frequency of training [/week]	$R_s = 0.31$	$R_s = 0.20$	$R_s = -0.03$	$R_s = 0.13$
	$P < 0.001$	$P = 0.02$	$P = 0.67$	$P = 0.13$
Change in training volume [h/week]	$R_s = 0.31$	$R_s = 0.13$	$R_s = 0.04$	$R_s = 0.14$
	$P < 0.001$	$P = 0.11$	$P = 0.68$	$P = 0.12$
Change in time spent in > 90% of HRmax [min]	$R_s = 0.26$	$R_s = 0.09$	$R_s = -0.05$	$R_s = 0.11$
	$P = 0.003$	$P = 0.26$	$P = 0.58$	$P = 0.19$
Change in time spent in 80–90% of HRmax [min]	$R_s = 0.27$	$R_s = 0.12$	$R_s = 0.02$	$R_s = 0.17$
	$P = 0.003$	$P = 0.16$	$P = 0.80$	$P = 0.04$
Change in TSS [average/week]*	$R_s = 0.00$	$R_s = -0.16$	$R_s = -0.22$	$R_s = 0.06$
	$P = 0.98$	$P = 0.26$	$P = 0.11$	$P = 0.66$
Change in TSS [maximum/week]*	$R_s = -0.02$	$R_s = -0.15$	$R_s = -0.24$	$R_s = -0.13$
	$P = 0.48$	$P = 0.28$	$P = 0.09$	$P = 0.40/$
VO ₂ max [ml/kg/min]	$R_s = 0.03$	$R_s = 0.05$	$R_s = -0.18$	$R_s = -0.03$
	$P = 0.81$	$P = 0.70$	$P = 0.21$	$P = 0.80$
Training experience [years]	$R_s = 0.10$	$R_s = 0.11$	$R_s = 0.15$	$R_s = 0.13$
	$P = 0.25$	$P = 0.18$	$P = 0.07$	$P = 0.12$
BMI [kg/m ²]	$R_s = 0.00$	$R_s = 0.00$	$R_s = 0.04$	$R_s = 0.05$
	$P = 0.96$	$P = 0.98$	$P = 0.61$	$P = 0.95$
Age [years]	$R_s = 0.07$	$R_s = 0.01$	$R_s = 0.04$	$R_s = 0.07$
	$P = 0.39$	$P = 0.61$	$P = 0.60$	$P = 0.37$

*Data available for 47/130 cyclists.

require a 3–6-month period of complete rest [40]. As shown in our study, the COVID-19-positive cyclists decreased their training intensity temporarily. To clarify, their return to training due to lack of information was not supervised as part of this research and could include cardiologic examination, troponins, MRI, and ECG, but we cannot exclude independent decisions to return to physical activity. Many athletes with confirmed COVID-19 on PCR testing are minimally symptomatic or asymptomatic. Return to outdoor exercise 7 days or later after two negative SARS-CoV-2 tests carries a very low risk of infecting others. According to guidelines, including omicron variants, routine medical investigation for athletes, including cardiac screening, is not recommended in those returning to exercise following COVID-19. The protocol for return to exercise for recreational and elite athletes assumes a stepwise (50%/75%/100%) increase of training intensity every 3 days [41].

There are a number of studies reporting on the effect of the COVID-19 pandemic on mental health and quality of life in the general population [42]. It was reported that 37% experience post-traumatic stress symptoms, 17% depression, 21% anxiety, 7% insomnia, while 22% suffer an increased stress level [43]. One of the present

study's unexpected observations was the higher physical, social relation, and environmental domain results in WHOQOL scoring among cyclists with a history of SARS-CoV-2 infection, but not in the psychological domain. The psychological domain was the only one that did not increase. We hypothesize that cyclists are highly individual athletes with good self-motivation, thus retaining a good psychological state with the experience of mild symptoms of COVID-19, producing the subjective evaluation of other domains: "I'm strong, nothing bad happened to me". A similar tendency was observed when cyclists were asked whether they knew a person who had died from COVID-19 or not. In this case, the psychological domain was also less favorable, with a lower score in athletes who had experienced bereavement, while other domains did not change. General norms for the WHOQOL-BREF domains were found to be higher than in our study: in non-pandemic conditions, 73.5 (SD = 18.1) for the physical health domain, 70.6 (14.0) for psychological well-being, 71.5 (18.2) for social relationships, and 75.1 (13.0) for the environmental domain [44]. We could speculate that higher scores in the somatic, social and environmental Quality of Life questionnaire in those cyclists that were infected by SARS-CoV-2

as compared to those that were not infected could represent a reduction in pandemic fear. This is opposite to the lower quality of life described in the literature after mild and severe disease courses. Among US professional endurance athletes, more than 60% reported the negative influence of the pandemic on training motivation, and 28% felt depressed. However, the authors did not analyze the impact of past SARS-CoV-2 infection on the eventual resolution of these symptoms [31]. In Italian elite athletes, self-determined motivation during COVID-19 was higher in individual than team athletes, elite vs. non-elite, and females vs. males. A significant positive relationship was observed between distress and mental health and intolerance of uncertainty and physical health [19]. The results obtained among cyclists who recovered from SARS-CoV-2 infection suggest that it is an important factor contributing to the quality of life of athletes under stressful and uncertain situations.

During lockdown in Austria, the psychological domain was observed to be similar and correlated positively with higher-income work activity [45]. The WHOQOL-BREF psychological domain was strongly dependent on physical activity; the authors observed mean scores of 62.8, 64.1, 69.1, 70.3, and 73.2 in cases with 0, 1, 2, 3 and 4 days of physical activity, respectively [45]. Our study's overall mean psychological domain score was 66.2 ± 12.2 and was also positively correlated with training frequency ($R_s = 0.20$, $p = 0.02$). Interestingly though, we observed a stronger positive correlation between training frequency, volume, and intensity with the physical domain than with psychological status. The association between physical activity and mental health, as found in our study, is well established. Increased duration and greater intensity of physical activity were both associated with a further reduction in the prevalence of depression in men [46].

In a study of physically active pregnant women in the second trimester, the WHOQOL psychological, social, and environmental domains did not correlate with the intensity of physical activity and volume. In the physical domain, a weak ($R_s = 0.16$) correlation between WHOQOL and highest intensity activity was observed [47]. It could also be speculated that a lower social domain score among cyclists resulted from isolation from general society to concentrate on healthy living and maintaining year-to-year goals in training. In this situation, the dominant role of training patterns should be emphasized in the physical domain, which by the way kept them also relatively psychologically healthy.

The study limitations should also be stressed. Firstly, the research encompassed a group of cyclists in Poland, and caution needs to be taken in extrapolating the findings to groups of such ath-

letes in other countries, as the epidemiological situation, imposed restrictions, and available resources may differ. Secondly, the study employed an anonymous online questionnaire, which was an appropriate approach during the pandemic and enabled a relatively high number of participants to be reached. Additional limitations are the incomplete availability of the TSS score and VO_2 max and the online sampling method.

In conclusion, cyclists maintained their volume and frequency of training during the pandemic compared to 2019. Still, they reduced the intensity of training by lowering the time spent on high-intensity intervals. Also, changes in the pattern were more likely to have been affected by their physical status rather than by their psychogenic condition. Moreover, the training pattern of cyclists who experienced death of family members or friends due to COVID-19 remained unaffected; however, they obtained lower psychological domain scores in WHOQOL.

Funding

This research was partially covered by the funds from the project entitled "Rozwój Uniwersyteckiego Centrum Badań Sportowo-Medycznych w Poznaniu" number NdS/544750/2021/2022 implemented within the program "Nauka dla Społeczeństwa" of the Ministry of Education and Science (Poland).

Ethical approval

The study was performed between 16 and 30 November 2020 and approved by the Bioethics Committee of Poznan University of Medical Sciences (KB – 170/21).

Conflict of interest

The authors declare no conflict of interest.

References

1. Rzymiski P, Nowicki M. COVID-19-related prejudice toward Asian medical students: a consequence of SARS-CoV-2 fears in Poland. *J Infect Public Health* 2020; 13: 873-76.
2. Betancourt JA, Rosenberg MA, Zevallos A, Brown JR, Mileski M. The impact of COVID-19 on telemedicine utilization across multiple service lines in the United States. *Healthcare (Basel)* 2020; 8: 380.
3. Papandreou C, Arija V, Aretouli E, Tsilidis KK, Bulló M. Comparing eating behaviours, and symptoms of depression and anxiety between Spain and Greece during the COVID-19 outbreak: cross-sectional analysis of two different confinement strategies. *Eur Eat Disord Rev* 2020; 28: 836-46.
4. Bendau A, Plag J, Kunas S, Wyka S, Strohle A, Petzold MB. Longitudinal changes in anxiety and psychological distress, and associated risk and protective factors during

- the first three months of the COVID-19 pandemic in Germany. *Brain Behav* 2020; 11: e01964.
5. Sidor A, Rzymiski P. Dietary choices and habits during COVID-19 lockdown: experience from Poland. *Nutrients* 2020; 12: 1657.
 6. Zhang J, Zhang Y, Huo S, et al. Emotional eating in pregnant women during the COVID-19 pandemic and its association with dietary intake and gestational weight gain. *Nutrients* 2020; 12: 2250.
 7. Kriaucioniene V, Bagdonaviciene L, Rodríguez-Pérez C, Petkeviciene JJN. Associations between changes in health behaviours and body weight during the COVID-19 quarantine in Lithuania: the Lithuanian COVIDiet Study. *Nutrients* 2020; 12: 3119.
 8. Scarano A, Inchingolo F, Rapone B, Festa F, Tari SR, Lorusso F. Protective face masks: effect on the oxygenation and heart rate status of oral surgeons during surgery. *Int J Environ Res Public Health* 2021; 18: 2363.
 9. Scarano A, Inchingolo F, Lorusso F. Environmental disinfection of a dental clinic during the Covid-19 pandemic: a narrative insight. *Biomed Res Int* 2020; 2020: 8896812.
 10. Reddy RK, Charles WN, Sklavounos A, Dutt A, Seed PT, Khajuria A. The effect of smoking on COVID-19 severity: a systematic review and meta-analysis. *J Med Virol* 2021; 93: 1045-56.
 11. Pollard MS, Tucker JS, Green HD. Changes in adult alcohol use and consequences during the COVID-19 pandemic in the US. *JAMA Netw Open* 2020; 3: e2022942.
 12. Jukic I, Calleja-González J, Cos F, et al. Strategies and solutions for team sports athletes in isolation due to COVID-19. *Sports (Basel)* 2020; 8: 56.
 13. Nhamo G, Dube K, Chikodzi D. Impact of COVID-19 on the Global Sporting Industry and Related Tourism, in Counting the Cost of COVID-19 on the Global Tourism Industry. Springer International Publishing Switzerland 2020.
 14. Walsh NP. Recommendations to maintain immune health in athletes. *Eur J Sport Sci* 2018; 18: 820-31.
 15. Diment BC, Fortes MB, Edwards JP, et al. Exercise intensity and duration effects on in vivo immunity. *Med Sci Sports Exerc* 2015; 47: 1390-98.
 16. Walsh NP, Oliver SJ. Exercise, immune function and respiratory infection: an update on the influence of training and environmental stress. *Immunol Cell Biol* 2016; 94: 132-9.
 17. Gleeson M, Pyne DB, Elkington LJ, et al. Developing a multi-component immune model for evaluating the risk of respiratory illness in athletes. *Exerc Immunol Rev* 2017; 23: 52-64.
 18. Edwards JP, Walsh NP, Diment PC, Roberts R. Anxiety and perceived psychological stress play an important role in the immune response after exercise. *Exerc Immunol Rev* 2018; 24: 26-34.
 19. Casali N, Cerea S, Moro T, Paoli A, Ghisi M. Just do it: high intensity physical activity preserves menatl and physical health in elite and non-elite athletes durign COVID-19. *Front Physiol* 2021; 12: 757150.
 20. Rzymiski P, Burzyński B, Knapik M, Kociszewski J, Wilczak M. How to balance the treatment of stress urinary incontinence among female athletes? *Arch Med Sci* 2021; 17: 314-22.
 21. Cochran WG. Sampling techniques. Wiley, USA 1977.
 22. Geldsetzer P. Use of rapid online surveys to assess people's perceptions during infectious disease outbreaks: a cross-sectional survey on COVID-19. *J Med Internet Res* 2020; 22: e18790.
 23. Coggan AR. Training and racing using a power meter: an introduction. Level II Coaching Manual. USA: Colorado Springs, 2003.
 24. WHOQOL Group. Development of the World Health Organization WHOQOL-BREF quality of life assessment. The WHOQOL Group. *Psychol Med* 1998; 28: 551-8.
 25. Skevington SM, Lotfy M, O'Connell KA. The World Health Organization's WHOQOL-BREF quality of life assessment: psychometric properties and results of the international field trial. A report from the WHOQOL group. *Qual Life Res* 2004; 13: 299-310.
 26. Lewko J, Misiak B, Sierżantowicz R. The relationship between mental health and the quality of life of Polish nurses with many years of experience in the profession: a cross-sectional study. *Int J Environ Res Public Health* 2019; 16: 1798.
 27. World Health Organization. WHOQOL-BREF: introduction, administration, scoring and generic version of the assessment: field trial version. World Health Organization 1996.
 28. Astorino TA, Edmunds RM, Clark A, et al. High-Intensity interval training increases cardiac output and VO2max. *Med Sci Sports Exerc* 2017; 49: 265-73.
 29. Klika RJ, Alderdice MS, Kvale JJ, Kearney JT. Efficacy of cycling training based on a power field test. *J Strength Cond Res* 2007; 21: 265-69.
 30. Naves JPA, Viana RB, Rebelo ACS, et al. Effects of high-intensity interval training vs. sprint interval training on anthropometric measures and cardiorespiratory fitness in healthy young women. *Front Physiol* 2018; 9: 1738.
 31. Roche M, Sainani K, Noordsey D, Fredericson M. The impacts od COVID-19 on mental health and training in US professional endurance athletes. *Clin J Sport Med* 2021; 32: 290-6.
 32. Chan ZYS, Peeters R, Cheing G, Ferber R, Cheung RTH. Evaluation of COVID-19 restrictions on distance runners' training habits using wearable trackers. *Front Sport Active Living* 2022; 3: 812214.
 33. Muriel X, Courel-Ibáñez J, Cerezuela-Espejo V, Pallarés JG. Training load and performance impairments in professional cyclists during COVID-19 lockdown. *Int J Sports Physiol Perform* 2021; 16: 735-8.
 34. Khalid Z, Farheen H, Tariq MI, Amjad I. Effectiveness of resistance interval training versus aerobic interval training on peak oxygen uptake in patients with myocardial infarction. *J Pak Med Assoc* 2019; 69: 1194-8.
 35. Hong KH, Choi JP, Hong SH, et al. Predictors of mortality in Middle East respiratory syndrome (MERS). *Thorax* 2018; 73: 286-9.
 36. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; 395: 1054-62.
 37. Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020; 368: m1091.
 38. Mitrani RD, Dabas N, Goldberger JJ. COVID-19 cardiac injury: implications for long-term surveillance and outcomes in survivors. *Heart Rhythm* 2020; 17: 1984-90.
 39. Wilson MG, Hull JH, Rogers J, et al. Cardiorespiratory considerations for return-to-play in elite athletes after COVID-19 infection: a practical guide for sport and exercise medicine physicians. *Br J Sports Med* 2020; 54: 1157-61.
 40. Barker-Davies RM, O'Sullivan O, Senaratne KPP, et al. The Stanford Hall consensus statement for post-COVID-19 rehabilitation. *Br J Sports Med* 2020; 54: 949-59.

41. Hughes DC, Orchard JW, Partridge EM, La Gerche A, Broderick C. Return to exercise post-COVID-19 infection: a pragmatic approach in mid-2022. *J Sci Med Sport* 2022; 25: 544-7.
42. dos Santos CF, Picó-Pérez M, Morgado P. COVID-19 and mental health – what do we know so far? *Front Psychiatry* 2020; 11: 565698.
43. Rossi R, Socci V, Talevi D, et al. COVID-19 pandemic and lockdown measures impact on mental health among the general population in Italy. *Front Psychiatry* 2020; 11: 790.
44. Hawthorne G, Herrman H, Murphy B. Interpreting the WHOQOL-Brèf: preliminary population norms and effect sizes. *Soc Indic Res* 2006; 77: 37-59.
45. Pieh C, Budimir S, Probst T. The effect of age, gender, income, work, and physical activity on mental health during coronavirus disease (COVID-19) lockdown in Austria. *J Psychosom Res* 2020; 136: 110186.
46. Currier D, Lindner R, Spittal MJ, Cvetkovski S, Pirkis J, English DR. Physical activity and depression in men: Increased activity duration and intensity associated with lower likelihood of current depression. *J Affect Disord* 2020; 260: 426-31.
47. Krzepota J, Sadowska D, Biernat E. Relationships between physical activity and quality of life in pregnant women in the second and third trimester. *Int J Environ Res* 2018; 15: 2745.