

Electrocardiographic repolarization parameters in children with arrhythmias and excessive body weight

Joanna Jaromin¹, Grażyna Markiewicz-Łoskot¹, Lesław Szydłowski², Agnieszka Kulawik³,
Agnieszka Chlebowczyk¹, Ewa Moric-Janiszewska⁴

¹Department of Nursing and Social Medical Problems, Faculty of Health Sciences, Medical University of Silesia, Katowice, Poland

²Department of Pediatric Cardiology, Faculty of Medical Sciences, Medical University in Silesia, Katowice, Poland

³Faculty of Science and Technology, University of Silesia, Katowice, Poland

⁴Department of Biochemistry, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia in Katowice, Poland

Submitted: 11 July 2023; **Accepted:** 9 December 2023

Online publication: 1 August 2024

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/176675>

Copyright © 2024 Termedia & Banach

Corresponding author:

Dr. Joanna Jaromin
Department of Nursing
and Social
Medical Problems
Faculty of
Health Sciences
Medical University
of Silesia
Katowice, Poland
E-mail: jjaromin@sum.edu.pl

Abstract

Introduction: In clinical diagnosis, electrocardiographic repolarization abnormalities with prolongation of QT and especially TpTe intervals are important for assessing the risk of malignant ventricular arrhythmias. The aim of the study was to compare the electrocardiological values of repolarization parameters in children with and without arrhythmias, according to body weight.

Material and methods: Repolarization parameters were compared in a group of children with excessive and normal body weight (BMI, Cole's index) with ventricular and supraventricular arrhythmias and in another group of children (matched for age and sex, with no arrhythmia and with normal weight) as the control group. The repolarization parameters TpTe, QTp and QT interval corrected Bazett (QTcB) and Fridericia (QTcF) formulas were measured in lead V5 of the 12-lead ECG.

Results: The durations of TpTe and QTcF intervals were significantly longer in children with ventricular arrhythmias with abnormal body weight compared to children with supraventricular arrhythmias with abnormal body weight. Comparing children with normal weight between the ventricular and supraventricular groups, significant prolongation of only the TpTe interval was observed. Moreover, there were statistically significant differences ($p < 0.0001$) in the TpTe interval among the 5 groups: children with ventricular and supraventricular arrhythmia with abnormal and normal weight, as well as the control group depending on BMI and Cole's index.

Conclusions: In children with ventricular arrhythmia, only the TpTe parameter was significantly longer compared to children with supraventricular arrhythmia and children without arrhythmia, regardless of body weight. In children with arrhythmia, regardless of body weight, the TpTe parameter should be measured in addition to routine QT. Further studies on a larger scale are needed to assess the clinical importance of the TpTe interval in children with arrhythmias and excessive body weight.

Key words: children, body weight, electrocardiography, cardiac arrhythmias.

Introduction

Overweight and obesity in children are considered among the most significant global public health problems [1–3]. Obesity at a young age

is considered to be a risk factor for cardiovascular morbidity and mortality [4]. Abnormal weight in early life predisposes not only to ischemic heart disease, but also to arrhythmias [5, 6]. Ventricular or supraventricular arrhythmia may occur sporadically, intermittently or permanently in children at any age from the fetal age [7]. Cardiac arrhythmias often resolve spontaneously during the child's development, while others do not manifest any clinical symptoms, but may cause the development of cardiomyopathy or may develop into life-threatening arrhythmias and lead to sudden cardiac death [8–11]. Monitoring and analysis of a resting ECG and/or a 24-hour Holter ECG recording with recorded arrhythmia is essential in the diagnosis of arrhythmias as a potential cause of syncope, circulatory failure, as well as in assessing the risk of sudden cardiac death [12–15]. Changes in the period of ventricular repolarization – QT interval and TpTe (Tpeak-Tend) interval – allow the assessment of electrical instability of the heart muscle. In clinical diagnosis, electrocardiographic repolarization abnormalities with prolongation of QT and TpTe intervals (Tpeak-Tend) are important for assessing the risk of malignant ventricular arrhythmias. In the literature reports, the TpTe repolarization parameter seems to be a sensitive indicator of arrhythmogenesis [16–21]. There are few studies in the literature that have analyzed the ECG in obese children, and there are no studies involving children with arrhythmias taking into account the body mass index. The aim of the study was to compare the electrocardiological values of repolarization parameters in children with and without arrhythmias, depending on body weight.

Material and methods

Study data were obtained through a retrospective analysis of medical records of children hospitalized in the Department of Pediatric Cardiology of the Saint John Paul II Upper Silesian Child Health Centre in Katowice. Repolarization parameters were compared in children with excessive and normal body weight (assessed by BMI and Cole's index) with ventricular and supraventricular arrhythmias (study group $N = 88$) and in children from the control group without diagnosed arrhythmias and with normal weight ($N = 34$). The repolarization parameters TpTe, QT peak and QT interval corrected Bazett (QTcB) and Fridericia (QTcF) formulas were measured in lead V5 of the 12-lead ECG. All anthropometric measurements were taken on the first day of hospitalization during the physical examination before starting the ECG.

When analyzing comparative characteristics, overweight and obese children were included in the group of children with 'abnormal body weight' (BMI index and Cole's index).

Studied subjects

The study group consisted of 88 children (38 girls and 50 boys) aged 4 to 18 years (mean age: 13 ± 5 years) with ventricular or supraventricular arrhythmias of unknown etiology (recorded in a standard ECG and in a 24-hour Holter ECG monitoring), and 34 children with chest pain admitted for diagnostic examinations excluding cardiovascular etiology served as a control group (22 girls and 12 boys) aged 4 to 18 years (mean age: 13 ± 3 years).

None of the study group subjects (including controls) had a positive personal or family history of sudden cardiac death or structural heart disease on physical examination (or) Doppler echocardiography or exercise arrhythmias (Bruce protocol).

In order to participate in this study, patients with arrhythmias had to fulfill all the following inclusion criteria: ventricular or supraventricular arrhythmias in a standard ECG and in a 24-hour Holter ECG, previous diagnosis of arrhythmias in anamnesis and in the physical examination, without episodes of sudden cardiac death in relatives of the child before the age of 30, without inflammation and ion disorders (K, Ca, Mg), no use of drugs that change the repolarization period, and no use of stimulants (cigarettes, alcohol). For patients in the control group, the inclusion criteria were: no history of symptoms suggesting arrhythmias, no arrhythmias in the ECG and 24-hour Holter ECG, normal physical examination result, with normal echocardiographic results, normal laboratory test results excluding inflammation and ionic disturbances (K, Ca, Mg), negative family history of structural and electrophysiological abnormalities of the heart, no episodes of sudden cardiac death in the child's relatives before the age of 30 years, no use of drugs that change the repolarization period, no use of stimulants (cigarettes, alcohol).

Anthropometric measurements

Body weight and height were assessed for all children in the present study. The children were tested in underwear and without shoes. Each measurement was taken as the average of three consecutive measurements. In children, body weight and height were measured in three repetitions. Body weight was measured with a Charder MS6110 medical scale with a height gauge HM202P (JAWAG, Morawica, Poland) with an accuracy of ± 100 g. Children's height was measured with a stadiometer attached to the scale with an accuracy of ± 0.1 cm. Mean values of height and weight were obtained from the three measurements to calculate the body mass index (BMI = body weight in kg/height in m^2), which was ad-

justed to the standards published by Cole *et al.* and Cole's index (LMS) ($(\% = \text{body weight} \times (\text{standard body length})^2 / \text{standard body weight} \times (\text{body length})^2 \times 100)$) [22].

The children were divided according to their nutritional status, once in reference to BMI and the second to Cole's index. According to BMI, underweight was defined as children with a value below the 5th percentile, slimness between the 5th and 25th percentile and obese as a value above the 95th percentile. Taking into account Cole's index, it was assumed that children with the index value below 75% are characterized by wasting, and children with the index value > 110% are overweight [23].

Electrocardiography

Standard ECG recordings were made in the supine position on the recorder model AT2 plus Schiller AG (Baar, Switzerland). Measurements were made manually in the fifth precordial lead (V5) of the 12-lead ECG. The analyzed measurements were the average of three consecutive QRS-T evolutions with a paper travel of 50 mm/s and a standard feature amplitude of 1 mV = 1 cm. In the analysis of the repolarization period, the duration of individual repolarization parameters and the amplitude were assessed. From three consecutive QRS-T evolutions, RR intervals, total repolarization time (QT interval) and TpTe interval (Tpeak-Tend) were determined and recorded under magnification with a magnifying glass using a distance stepper. The total period of QT repolarization was determined from the beginning of the Q wave to the end of the T wave, defined as the point of return of the descending arm of the T wave to the isoelectric line, excluding the U wave [24, 25]. Corrected QT duration for heart rate (QTcB) was calculated using the Bazett (QT/\sqrt{RR}) and Fridericia ($QT/RR^{1/3}$) formula [26, 27]. In the case of two T-wave peaks, the first T-wave peak was taken into account in the measurements [28–30]. All measurements were analyzed blindly by two independent investigators with uniformity tests without access to the results obtained and clinical data. The highest quality ECG was analyzed without network interference with the exception of the flat T wave. 24-hour Holter was performed in all children ($n = 122$).

Statistical analysis

The calculations and graphs used in the article (except for minor modifications) were made in Statistica software (version 13.1, StatSoft Poland). In the statistical analysis tests for two independent samples and a test for five independent groups were used. The proposed tests depend on the normality of the distribution of the studied character-

istic in the considered groups. If the test of normality of distribution in the groups under study did not show statistically significant differences with the normal distribution, a parametric test (the *t*-test for two independent samples) was used. Otherwise, non-parametric tests (Mann-Whitney *U* test or Kruskal-Wallis test) were used. The results are presented in the corresponding tables. All tables include the arithmetic mean and the standard deviation for each sample, as well as the *p*-value for a given test. A significance level of 0.05 was adopted in the statistical analysis. For *p*-values less than 0.05, statistically significant differences in the study populations can be inferred.

Results

Most children had normal weight (ventricular arrhythmias – $n = 33$, 71.7% (BMI), $n = 24$, 52.1% (Cole's index), supraventricular arrhythmias – $n = 29$, 69% (BMI), $n = 27$, 64.2% (Cole's index)). According to Cole's index, 37 (30.3%) children with arrhythmias were overweight, while the BMI index indicated overweight in 22 (18%) children of this group. All children in the control group had normal weight (Table I).

The duration of the TpTe, QTcB and QTcF intervals were significantly longer in the group of children with arrhythmias and with abnormal weight compared to the values of the TpTe, QTcB, and QTcF intervals of the control group with normal weight according to the BMI index (Table II).

The durations of TpTe, QTcB and QTcF intervals were significantly longer in the group of children with arrhythmias and with abnormal weight compared to the values of the TpTe, QTcB, and QTcF intervals of the control group with normal weight according to Cole's index (Table III).

Only the duration of the TpTe interval was significantly longer in the group of children with ventricular arrhythmias and with abnormal weight compared to the value of the TpTe interval of the group of children with supraventricular arrhythmias and abnormal weight according to the BMI index (Table IV).

The durations of TpTe and QTcF intervals were significantly longer in the group of children with ventricular arrhythmias and with abnormal weight compared to the values of the TpTe and QTcF intervals of the group of children with supraventricular arrhythmias and with abnormal weight according to Cole's index (Table V).

Only the duration of the TpTe interval was significantly longer in the group of children with ventricular arrhythmias and with normal weight compared to the value of the TpTe interval of the group of children with supraventricular arrhythmias and normal weight according to the BMI index (Table VI).

Table I. Nutritional status in children with ventricular and supraventricular arrhythmias according to weight assessed by body mass index (BMI) and Cole's index

Parameter	Total cohort (n = 122)	Children with ventricular arrhythmias (n = 46)	Children with supraventricular arrhythmias (n = 42)	Control group (n = 34)
Age (median)	13.4	13.6	13.4	13.3
Female, n (%)	60 (49)	21 (46)	17 (40)	22 (65)
Male, n (%)	62 (51)	25 (54)	25 (60)	12 (35)
Nutritional status – BMI index:				
Underweight (< 5 th percentile), n (%)	0	0	0	0
Normal weight (25 th –85 th percentile), n (%)	96 (78.6)	33 (71.7)	29 (69)	34 (100)
Overweight (85 th –95 th percentile), n (%)	22 (18)	9 (19.5)	13 (30.9)	0
Obese (> 95 th percentile), n (%)	4 (3.2)	4 (8.6)	0	0
Nutritional status – Cole's index:				
Emaciation (< 75%), n (%)	0	0	0	0
Normal weight (90–110%), n (%)	85 (69.6)	24 (52.1)	27 (64.2)	34 (100)
Overweight (> 110%), n (%)	37 (30.3)	22 (47.8)	15 (35.7)	0

Table II. Characteristics of QT, QTp, TpTe, QTcB and QTcF intervals in the group of children with arrhythmias with abnormal weight and children from the control group with normal weight according to BMI index

Parameter [ms]	Children with arrhythmias		Children from the control group		Test	P-value
	Abnormal weight (n = 26)		Normal weight (n = 34)			
	Mean	Standard deviation	Mean	Standard deviation		
QT	361.9	22.3	357.6	24.9	Mann-Whitney U test	0.338
QTp	278.8	21.4	286.8	24.6	Mann-Whitney U test	0.178
TpTe	83.1	4.7	70.9	4.5	Mann-Whitney U test	0.000
QTcB	406.5	23.8	391.1	21.4	Student's t-test	0.010
QTcF	390.8	17.6	379.4	18.1	Student's t-test	0.017

Table III. Characteristics of QT, QTp, TpTe, QTcB and QTcF intervals in the group of children with arrhythmias with abnormal weight and children from the control group with normal weight according to Cole's index

Parameter [ms]	Children with arrhythmias		Children from the control group		Test	P-value
	Abnormal weight (n = 37)		Normal weight (n = 34)			
	Mean	Standard deviation	Mean	Standard deviation		
QT	365.9	22.9	357.6	24.9	Mann-Whitney U test	0.079
QTp	281.6	22.3	286.8	24.6	Mann-Whitney U test	0.338
TpTe	84.3	5.5	70.9	4.5	Mann-Whitney U test	0.000
QTcB	408.8	24.0	391.1	21.4	Student's t-test	0.001
QTcF	393.7	17.2	379.4	18.1	Student's t-test	0.001

Table IV. Characteristics of QT, QTp, TpTe, QTcB and QTcF intervals in the group of children with ventricular arrhythmias with abnormal weight and children with supraventricular arrhythmias with abnormal weight according to BMI index

Parameter [ms]	Children with arrhythmias				Test	P-value
	Children with ventricular arrhythmias and abnormal weight (n = 13)		Children with supraventricular arrhythmias and abnormal weight (n = 13)			
	Mean	Standard deviation	Mean	Standard deviation		
QT	366.2	21.0	357.7	23.5	Student's <i>t</i> -test	0.343
QTp	280.8	19.8	276.9	23.6	Student's <i>t</i> -test	0.656
TpTe	85.4	5.2	80.8	2.8	Mann-Whitney <i>U</i> test	0.044
QTcB	411.8	23.5	401.2	23.8	Student's <i>t</i> -test	0.262
QTcF	395.6	14.4	385.9	19.8	Student's <i>t</i> -test	0.164

Table V. Characteristics of QT, QTp, TpTe, QTcB and QTcF intervals in the group of children with ventricular arrhythmias with abnormal weight and children with supraventricular arrhythmias with abnormal weight according to Cole's index

Parameter [ms]	Children with arrhythmias				Test	P-value
	Children with ventricular arrhythmias and abnormal weight (n = 22)		Children with supraventricular arrhythmias and abnormal weight (n = 15)			
	Mean	Standard deviation	Mean	Standard deviation		
QT	370.0	22.7	360.0	22.7	Student's <i>t</i> -test	0.196
QTp	283.2	22.3	279.3	22.8	Mann-Whitney <i>U</i> test	0.766
TpTe	86.8	5.7	80.7	2.6	Mann-Whitney <i>U</i> test	0.000
QTcB	414.5	22.3	400.4	24.8	Student's <i>t</i> -test	0.079
QTcF	398.7	13.2	386.3	19.9	Student's <i>t</i> -test	0.027

Table VI. Characteristics of QT, QTp, TpTe, QTcB and QTcF intervals in the group of children with ventricular arrhythmias with normal weight and children with supraventricular arrhythmias with normal weight according to BMI index

Parameter [ms]	Children with arrhythmias				Test	P-value
	Children with ventricular arrhythmias and normal weight (n = 33)		Children with supraventricular arrhythmias and normal weight (n = 29)			
	Mean	Standard deviation	Mean	Standard deviation		
QT	371.5	22.9	363.4	26.2	Mann-Whitney <i>U</i> test	0.432
QTp	282.7	24.1	283.4	26.2	Mann-Whitney <i>U</i> test	0.500
TpTe	88.5	6.2	80.0	3.8	Mann-Whitney <i>U</i> test	< 0.001
QTcB	412.3	27.2	402.4	27.8	Student's <i>t</i> -test	0.163
QTcF	397.9	19.7	388.7	22.2	Student's <i>t</i> -test	0.088

Table VII. Characteristics of QT, QTp, TpTe, QTcB and QTcF intervals in the group of children with ventricular arrhythmias with normal weight and children with supraventricular arrhythmias with normal weight according to Cole’s index

Parameter [ms]	Children with arrhythmias				Test	P-value
	Children with ventricular arrhythmias and normal weight (n = 24)		Children with supraventricular arrhythmias and normal weight (n = 27)			
	Mean	Standard deviation	Mean	Standard deviation		
QT	370.0	22.5	362.6	27.0	Student’s t-test	0.295
QTp	281.3	23.6	282.6	27.0	Student’s t-test	0.851
TpTe	88.3	6.4	80.0	3.9	Mann-Whitney U test	0.000
QTcB	410.0	29.3	402.9	27.6	Student’s t-test	0.378
QTcF	395.9	22.1	388.7	22.3	Student’s t-test	0.251

Only the duration of the TpTe interval was significantly longer in the group of children with ventricular arrhythmias and with normal weight compared to the value of the TpTe interval of the group of children with supraventricular arrhythmias and normal weight according to Cole’s index (Table VII).

There were no significant differences (TpTe interval: using BMI and Cole’s index) in the group of children with ventricular arrhythmias with abnormal body weight and children with the same arrhythmias with normal weight, as well as in the group with supraventricular arrhythmias in terms of weight in the TpTe interval (using BMI

and Cole’s index) and the QT, QTp, QTcB and QTcF intervals with their corrected values. Moreover, there were statistically significant differences in the TpTe interval between the 5 groups: children with ventricular and supraventricular arrhythmia with abnormal and normal weight, as well as the control group depending on BMI and Cole’s index (Figure 1).

Discussion

Compared to obese adult studies there are few studies on ECG changes in obese children. There are many studies focusing mainly on a group of young athletes to assess the risk of sudden car-




Figure 1. Statistically significant differences ($p < 0.0001$) in TpTe intervals among the 5 groups: children with ventricular and supraventricular arrhythmia with abnormal and normal weight, as well as control group (green color) depending on BMI index (blue color) and Cole’s index (red color). Values [ms; millisecond] are shown as the median, the quartiles and the range for each group

diac death, without reference to anthropometric measurements [31–35]. In clinical diagnosis, electrocardiographic repolarization abnormalities with prolongation of QT and TpTe intervals (Tpeak-Tend) are important for assessing the risk of malignant ventricular arrhythmias [16–21].

Our study showed a significantly longer duration of TpTe, QTcB and QTcF intervals in children with cardiac arrhythmias (ventricular or supraventricular) regardless of body weight (assessed by BMI and Cole's index) compared to healthy children. Similarly, previous studies by Sun *et al.* and Leotta *et al.* showed that the QTc interval was not associated with obesity in a healthy population of children [36, 37].

In the study of Yıldırım Yıldız *et al.*, involving 81 obese children, prolongation of the QT/QTc interval in the electrocardiogram at an early age was demonstrated in obese children [38]. Similarly, in studies by Paech *et al.*, a significant prolongation of the QTc and TpTe intervals was observed in obese healthy children compared to healthy lean children [39]. Radbill *et al.* observed that obesity was independently associated with the occurrence of postoperative arrhythmias in children with congenital heart defects [40]. Santini *et al.* presented different results, showing a significantly higher incidence of minor ECG abnormalities in underweight children and a lower incidence in obese children [41]. In subsequent studies by Paech *et al.*, involving 53 healthy obese children and 43 lean children from the control group, no differences between the groups in the ECG recording were found, and no ECG abnormalities were recorded in healthy obese children [42].

Kiess *et al.* noted that the results of research on the effect of obesity in children on the ECG recording so far are inconsistent. This may be due to the fact that the pathological changes in the ECG present in obese adult patients are not so far observable in childhood [43].

Our study showed significant prolongation of the TpTe interval (using BMI and Cole's index) and QTcF (only Cole's index) in children with ventricular arrhythmias with abnormal body weight compared to children with supraventricular arrhythmias with abnormal body weight (Tables IV and V). Comparing children with normal weight between the ventricular and supraventricular groups, significant prolongation of only the TpTe interval was observed (using the BMI and Cole's index) (Tables VI and VII).

In our study, it was also observed that Cole's index seems to be a more sensitive parameter than BMI, in accordance with a previous study by Abartes *et al.* [23].

The repolarization parameter TpTe interval is often assessed in the pediatric population [14, 16–21, 30, 44]. However, to our knowledge, there

are no scientific reports in the literature indicating a relationship between the parameters of the repolarization period in the group of children with cardiac arrhythmias (ventricular or supraventricular) and body weight (based on BMI and Cole's index). These are novel findings in obese children with cardiac arrhythmias, and no similar data have been found in available publications to date. Clinical usefulness of the TpTe in children with ventricular and supraventricular arrhythmias and excessive body weight seems to be an interesting issue. However, it requires further research, especially on larger groups of obese children with cardiac arrhythmias, taking into consideration the sex and age of patients.

In conclusion, in children with ventricular arrhythmia, only the TpTe parameter was significantly longer compared to children with supraventricular arrhythmia and children without arrhythmia, regardless of body weight. In children with arrhythmia, regardless of body weight, the TpTe parameter should be measured in addition to routine QT. Further studies on a larger scale are needed to assess the clinical importance of the TpTe interval in children with arrhythmias and excessive body weight.

Funding

No external funding.

Ethical approval

The study was approved by the Bioethics Committee of the Medical University of Silesia in Katowice (No. KNW/0022/KB/238/15).

Conflict of interest

The authors declare no conflict of interest.

References

1. Temiz F, Güneş H, Güneş H. Evaluation of atrial electromechanical delay in children with obesity. *Medicina* 2019; 55: 228.
2. Wong G, Srivastava G. Obesity management in children and adolescents. *Gastroenterol Clin N Am* 2023; 52: 443-55.
3. Jebile H, Kelly AS, O'Malley G, Baur LA. Obesity in children and adolescents: epidemiology, causes, assessment, and management. *Lancet Diabetes Endocrinol* 2022; 10: 351-65.
4. Alfaris N, Alqahtani A.M., Alamuddin N, Rigas G. Global impact of obesity. *Gastroenterol Clin N Am* 2023; 52: 277-93.
5. Singh GM, Danaei G, Farzadfar F, et al. The age-specific quantitative effects of metabolic risk factors on cardiovascular diseases and diabetes: a pooled analysis. *PLoS One* 2013; 8: e65174.
6. Sommer A, Twig G. The impact of childhood and adolescent obesity on cardiovascular risk in adulthood: a systematic review. *Curr Diab Rep* 2018; 18: 91.

7. Jaszczyszyn E, Panaszek B. Principles of electrophysiology, causes and classification of cardiac arrhythmias – prognostic and therapeutic implications. *Prim Care Rev* 2013; 15: 573-80.
8. Brignole M. Guidelines on management (diagnosis and treatment) of syncope – update 2004. The Task Force on Syncope, European Society of Cardiology. *Europace* 2004; 6: 467-537.
9. Wójcicka-Urbańska B, Kamińska H, Tomik A, et al. Arrhythmias in newborns – own experience. *Nowa Pediatrya* 2013; 4: 143-7.
10. Miszczak-Knecht M, Werner B. Benign arrhythmia in children. *N Ped* 2015; 19: 96-100.
11. Puranik R, Chow CK, Duflou JA, Kilborn MJ, McGuire MA. Sudden death in the young. *Heart Rhythm* 2005; 2: 1277-82.
12. Turska-Kmieć A, Ziótkowska L, Kawalec W. Modern concepts on the etiology of the idiopathic ventricular arrhythmias in children. *Prog Med* 2011; 12: 1046-53.
13. Yan GX, Antzelevitch C. Cellular basis for the normal T-wave and the electrocardiographic manifestation of the long QT syndrome. *Circulation* 1998; 98: 1928-36.
14. Markiewicz-Łoskot G, Kolarczyk E, Mazurek B, Łoskot M, Szydlowski L. Prolongation of electrocardiographic T wave parameters recorded during the head-up tilt table test as independent markers of syncope severity in children. *Int J Environ Res Public Health* 2020; 17: 6441.
15. Priori SG, Blomström-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *Eur Heart J* 2015; 36: 2793-867.
16. Pruszkowska-Skrzep P, Pluta S, Lenarczyk A, et al. A comparison of the clinical course of preexcitation syndrome in children and adolescents and in adults. *Cardiol J* 2007; 14: 384-90.
17. Kolarczyk E, Markiewicz-Łoskot G, Szydlowski L. The repolarization period during the head-up tilt test in children with vasovagal syncope. *Int J Environ Res Public Health* 2020; 17: 1908.
18. Antzelevitch C, Di Diego JM. Tpeak-Tend interval as a marker of arrhythmic risk. *Heart Rhythm* 2019; 16: 954-5.
19. Schäfer M, Frank BS, Ivy DD, et al. Repolarization dispersion is associated with diastolic electromechanical discoordination in children with pulmonary arterial hypertension. *J Am Heart Assoc* 2022; 11: e024787.
20. Bird K, Chan G, Lu H, et al. Assessment of hypertension using clinical electrocardiogram features: a first-ever review. *Front Med (Lausanne)* 2020; 7: 583331.
21. Jaromin J, Markiewicz-Łoskot G, Szydlowski L, Kulawik A. Diagnostic value of the TpTe interval in children with ventricular arrhythmias. *Int J Environ Res Public Health* 2021; 18: 12194.
22. Cole TJ, Flegal KM, Nicholls D, et al. Body mass index cut offs to define thinness in children and adolescents: international survey. *Br Med J* 2007; 335: 194.
23. Abrantes MM, Lamounier JA, Colosimo EA. Comparison of body mass index values proposed by Cole et al. (2000) and Must et al. (1991) for identifying obese children with weight-for-height index recommended by the World Health Organization. *Public Health Nutr* 2003; 6: 307-11.
24. Antzelevitch C, Sicouri S. Clinical relevance of cardiac arrhythmias generated by after depolarizations. Role of M cells in the generation of U waves, triggered activity and torsade de pointes. *J Am Coll Cardiol* 1994; 23: 259-77.
25. Haraguchi Y, Yosinaga M, Sarantuya J, et al. Interval representative of transmural dispersion of repolarization in children and young adolescents with congenital long QT syndrome. *Circ J* 2005; 69: 78-82.
26. Fridericia LS. Diesystolendauerim elektrokardiogramm bei normalen menschen und bei herzkranken. *Acta Med Scand* 1920; 53: 469-86.
27. Bazett .C. An analysis of time relations of electrocardiograms. *Heart* 1920; 7: 353-67.
28. Rautaharju PM, Surawicz B, Gettes LS. AHA/ACC/FHRS recommendations for the standardization and interpretation of the electrocardiogram: Part IV: The ST segment, T and U waves, and the QT interval: A scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society; endorsed by the International Society for Computerized Electrocardiology. *Circulation* 2009; 119: 241-50.
29. Takenaka K, Tomohiko A, Shimizu W, et al. Exercise stress test amplifies genotype-phenotype correlation in the LQT1 and LQT2 forms of the long-QT syndrome. *Circulation* 2003; 107: 838-44.
30. Markiewicz-Łoskot G. Electrocardiographic characteristics of a total of repolarization (QT), early repolarization phase (QTP) and late phase repolarization (TpTe) in healthy children and children with long QT syndrome. Medical University of Silesia: Katowice, Poland, 2009; Volume 8: 59–76.
31. Chandra N, Bastiaenen R, Papadakis M, et al. Prevalence of electrocardiographic anomalies in young individuals: relevance to a nationwide cardiac screening program. *J Am Coll Cardiol* 2014; 63: 2028-34.
32. De Lazzari C, Genuini I, Gatto MC, et al. Screening high school students in Italy for sudden cardiac death prevention by using a telecardiology device: a retrospective observational study. *Cardiol Young* 2017; 27: 74-81.
33. Bonaventura J, Rowin EJ, Maron MS, Maron BJ. Risks of the athletic field revisited: report of unusual occurrences of cardiac arrest and sudden death in professional soccer players. *Am J Med* 2023; 136: 315-21.
34. Sarto P, Zorzi A, Merlo Lk, et al. Serial versus single cardiovascular screening of adolescent athletes. *Circulation* 2021; 143: 1729-31.
35. Bohm P, Meyer T, Narayanan K, et al. Sports-related sudden cardiac arrest in young adults. *Europace* 2023; 25: 627-33.
36. Sun GZ, Li Y, Zhou XH, et al. Association between obesity and ECG variables in children and adolescents: a cross-sectional study. *Exp Ther Med* 2013; 6: 1455-62.
37. Leotta G, Maule S, Rabbia F, et al. Relationship between QT interval and cardiovascular risk factors in healthy young subjects. *J Hum Hypertens* 2005; 19: 623-7.
38. Yıldırım Yıldız N, Uçar T, Ramoğlu M, et al. Does obesity influence ventricular repolarisation in children? *Cardiol Young* 2021; 31: 568-76.
39. Paech C, Liebold A, Gebauer RA, et al. Relative QT interval prolongation and electrical inhomogeneity of cardiac repolarization in childhood obesity. *Progress Pediatr Cardiol* 2017; 47: 64-7.
40. Radbill AE, Smith AH, Van Driest SL, et al. Impact of obesity on post-operative arrhythmias after congenital heart surgery in children and young adults. *Cardiol Young* 2022; 32: 1820-5.
41. Santini M, Angela S, Fusco D, Colivicchi F, Gargaro A. Electrocardiographic characteristics, anthropometric

- features, and cardiovascular risk factors in a large cohort of adolescents. *EP Europace* 2018; 20: 1833-40.
42. Paech C, Anhalt M, Gebauer RA, et al. New normal limits for pediatric ECG in childhood obesity? Influence of childhood obesity on the ECG. *Progress Pediatr Cardiol* 2018; 48: 119-23.
 43. Kiess A, Körner A, Dähnert I, et al. Does obesity have an effect on the ECG in children? *J Pediatr Endocrinol Metab* 2020; 33: 585-9.
 44. Markiewicz-Łoskot G, Moric-Janiszewska E, Mazurek B, et al. Electrocardiographic T-wave parameters in families with long QT syndrome. *Adv Clin Exp Med* 2018; 27: 501-7.