

Association of cataract surgery with risk of dementia: a meta-analysis and systematic review of cohort studies with 720,075 participants

Qin Li¹, Liangqiong Zhu^{2*}

¹Department of Ophthalmology, Chongqing Hospital of Traditional Chinese Medicine, Chongqing, China

²Operation Room, Chongqing Hospital of Traditional Chinese Medicine, Chongqing, China

Submitted: 24 July 2024; **Accepted:** 14 September 2024

Online publication: 25 October 2024

Arch Med Sci 2024; 20 (5): 1720–1726

DOI: <https://doi.org/10.5114/aoms/193394>

Copyright © 2024 Termedia & Banach

***Corresponding author:**

Liangqiong Zhu

Operation Room

Chongqing Hospital

of Traditional

Chinese Medicine

Chongqing 400011

China

E-mail: 13883464995@163.

com

The treatment options for dementia patients remain limited in their efficacy. Consequently, there has been a growing research focus on identifying risk factors that could potentially prevent or delay the onset of dementia. Visual impairment has been recognized as a significant risk factor for dementia [1–3]. Cataract is the most prevalent reversible cause of blindness, and it is typically treated through surgical intervention to restore vision [4, 5]. Numerous studies [6–10] have investigated the association between cataract surgery and dementia. However, the impact of cataract surgery on the risk of developing Alzheimer's disease remains controversial. Therefore, we conducted a meta-analysis of existing literature on cataract surgery and its association with dementia, with the aim of providing a scientific foundation for strategies to prevent or delay the onset of Alzheimer's disease.

Methods. Search strategy The PubMed, Embase, and Web of Science databases from inception to March 31, 2024 were extensively searched for cohort studies on cataract surgery and dementia.

Inclusion criteria. (1). Study design: Cohort study; (2). Participant: Cataract patient; (3). Intervention: The observation group received cataract surgery, while the control group did not receive cataract surgery; (4) Outcomes: The primary outcome was all-cause dementia. Secondary outcomes were other types of dementia, including Alzheimer's disease, vascular dementia, etc.; (5). Effect size: relative risk (RR), odds ratio (OR), and hazard ratio (HR) with 95% confidence intervals (CI).

Exclusion criteria. (1) Non-English literature; (2) Review literature or case studies; (3) Data missing, or unable to extract, or the full text of the literature not available; (4) Republished literature.

Data extraction. When screening articles, we first read the title to eliminate irrelevant ones, and then further reviewed the abstract and full text to determine whether to include the article. The data collected were collated and analyzed independently by two researchers using Excel 2021. The extracted information included the first author, year of publication, region, data source, study design, sample size, type of dementia, cataract diagnosis, dementia diagnosis, follow-up time, age, outcome, effect size, 95% CI, and Newcastle-Ottawa Scale (NOS) score. We cross-examined the findings, and in the event of a dispute, discussed the resolution or consulted a third researcher.

Study quality assessment. NOS was used to evaluate the quality of included studies [11]. An NOS score ≥ 7 was considered to be of high quality.

Data synthesis and analysis. HR and 95%CI were calculated using Stata14.0 software. I^2 values were used to detect the heterogeneity between included studies. If $p > 0.1$ and $I^2 < 50\%$, there was no statistical heterogeneity, and a fixed effect model was used for analysis. If $p < 0.1$ and $I^2 \geq 50\%$, there was statistical heterogeneity, and a random effects model was used for analysis. Subgroup analyses were performed based on study design (retrospective vs. prospective), sample size ($< 10,000$ vs. $> 10,000$), region (Asian vs. non-Asian), and whether to adjust for APOE $\epsilon 4$ mutations (yes vs. no). The funnel plot method was used to analyze whether publication bias existed. Finally, sensitivity analysis was used to verify the stability of the results.

Results. Study characteristics. The number of items of literature preliminarily obtained through database retrieval was 462. After a thorough screening and careful review, five articles [6–10] were finally included for meta-analysis, involving 720,075 participants (Figure 1 A). The basic characteristics of the literature are presented in Table I.

Meta-analysis results. Meta-analysis showed that cataract surgery was not associated with a reduced risk of all-cause dementia (HR = 0.85,

95% CI (0.68, 1.06), $p = 0.158$) (Figure 1 B). However, it was associated with a decreased risk of Alzheimer’s disease (HR = 0.58, 95% CI (0.36, 0.93), $p = 0.025$) (Figure 1 B). Descriptive analysis revealed that cataract surgery did not correlate with a reduced risk of vascular dementia (HR = 1.03, 95% CI (0.51, 2.11), $p = 0.935$) (Figure 1 B), but it was associated with a lower risk of mild cognitive impairment (HR = 0.79, 95% CI (0.65, 0.97), $p = 0.023$) (Figure 1 B).

Subgroup analysis. Subgroup analysis was performed based on study design (retrospective vs. prospective). Subgroup analyses showed that cataract surgery was not associated with a reduced risk of all-cause dementia, regardless of whether the study design was prospective or retrospective (retrospective: HR = 0.91, 95% CI (0.64, 1.31), $p = 0.621$; prospective: HR = 0.80, 95% CI (0.57, 1.11), $p = 0.176$) (Figure 1 C).

Subgroup analysis was performed according to the study sample size ($< 10,000$ vs. $> 10,000$). Subgroup analyses showed that cataract surgery was not associated with a reduced risk of all-cause dementia, regardless of sample size $< 10,000$ or $> 10,000$ ($< 10,000$: HR = 0.89, 95% CI (0.54, 1.49), $p = 0.663$; $> 10,000$: HR = 0.84, 95% CI (0.61, 1.14), $p = 0.264$) (Figure 1 C).

Subgroup analysis was performed by region (Asian vs. non-Asian). Subgroup analyses showed




Figure 1. A – Preferred reporting items for systematic reviews and meta-analyses flowchart of included studies







Figure 1. Cont. **B** – Forest plot of association between cataract surgery and dementia risk; **C** – Forest plot of subgroup analysis of cataract surgery and the risk of all-cause dementia; **D** – Sensitivity analysis for the association between cataract surgery and the risk of all-cause dementia; **E** – Forest plot of subgroup analysis of cataract surgery and the risk of all-cause dementia (after excluding Lee et al. 2023)

Table 1. Basic characteristics of included studies

Author (year)	Country	Data source	Study design	Sample size	Population	Age	Definition of cataract	Diagnostic criteria for dementia	Follow-up period	Outcomes types	Confounders adjusted	NOS scores
Yu <i>et al.</i> 2015	China	National Health Insurance Research Database (NHIRD) of Taiwan	Retrospective cohort study	113123	Cataract surgery	79.4 ±4.9	ICD-9-CM code 366.x	ICD-9-CM codes 290.x or 331.0	10 years	ACD	Age, sex, CCI score, the interval between the first coding of cataract diagnosis and index date, hypertension, and diabetes	7
Miyata <i>et al.</i> 2018	Japan	The Fujiwara-kyo Eye Study	Prospective cohort study	2764	Cataract surgery	76.3 ±4.8	A Landolt ring chart, the slit-lamp examination	The Mini-Mental State Examination (MMSE)	NR	ACD, MCI	Age, sex, body mass index, education, BCVA, hypertension, diabetes, depression, and stroke	8
Ma <i>et al.</i> 2023	UK	UK Biobank	Prospective cohort study	300,823	Cataract surgery	40–69	ICD-9/ICD10	ICD-9/ICD-10	Mean follow-up of 8.4 years	ACD, AD, VD	Age, sex, education, APOE ε4, history of other types of eye surgery, unilateral or bilateral cataracts, cataract duration, ethnicity, body mass index, smoking status, alcohol use, and CHD	9
Lee <i>et al.</i> 2023	Korea	Korean National Health Insurance Service data	Retrospective cohort study	300327	Cataract surgery	66.79 ±8.90	ICD-10	ICD-10	6.08 ±3.06 years	ACD	Age, sex, income VA, smoking history, drinking habits, body mass index, diabetes, hypertension, dyslipidemia, stroke, depression, CHD, glaucoma, diabetic retinopathy, and age-related macular degeneration	8
Lee <i>et al.</i> 2022	USA	ACT study	Prospective cohort study	3038	Cataract surgery	73.8 ±6.0	CD-9, ICD-10	DSM-IV	Mean follow-up of 7.8 years	ACD, AD	Education, self-reported, White race, smoking, APOE ε4, sex, age groups at cataract diagnosis, diabetes, systolic blood pressure, hypertension, heart disease, cardiovascular disease, body mass index, self-rated health, CCI, number of activities of daily living limitations, at least 15 min of activity 3 times/week, performance-based physical function scores, Center For Epidemiologic Studies Depression Scale scores, retirement status, and self-reported difficulty with distance or near vision	9

ACD – all-cause dementia except AD and VD; AD – Alzheimer's disease, VD – vascular dementia, MCI – mild cognitive impairment, CCI – Charlson comorbidity index, BCVA – best correct visual acuity, CHD – coronary heart disease, VA – visual acuity, NOS – Newcastle-Ottawa Scale.

that cataract surgery was not associated with a reduced risk of all-cause dementia, regardless of region being Asian or non-Asian (Asian: HR = 0.84, 95% CI (0.61, 1.14), $p = 0.264$; non-Asian: HR = 0.89, 95% CI (0.54, 1.49), $p = 0.663$) (Figure 1 C).

Subgroup analyses were performed based on whether the included studies adjusted for APOE $\epsilon 4$ mutations (yes vs. no). Subgroup analyses showed that after adjusting for APOE $\epsilon 4$ mutations, cataract surgery was associated with a reduced risk of all-cause dementia (HR = 0.70, 95% CI (0.62, 0.80), $p < 0.0001$) (Figure 1 C). However, without adjusting for APOE $\epsilon 4$ mutations, cataract surgery was not associated with a reduced risk of all-cause dementia (HR = 0.98, 95% CI (0.71, 1.34), $p = 0.876$) (Figure 1 C).

Sensitivity analysis. The included literature was excluded item by item for sensitivity analysis. When the study by Lee *et al.* (2023) [6] was excluded, the meta-analysis results showed statistical significance (Figures 1 D, E).

Publication bias. As fewer than 10 articles were included, publication bias analysis was not possible.

Discussion. Our meta-analysis of existing studies showed that cataract surgery was not associated with a reduced risk of all-cause dementia, but it was associated with a decreased risk of developing Alzheimer's disease. The results of this meta-analysis are quite different from those of previous meta-analyses. Previous meta-analyses suggested that cataract surgery might help reduce the risk of all-cause dementia [12–14]. First, Liu *et al.*'s [12] meta-analysis pooled various types of dementia and found that cataract surgery was associated with a reduced risk of dementia. Additionally, a meta-analysis by Zhang *et al.* [13] found that cataract surgery was associated with a lower incidence of all-cause dementia and Alzheimer's disease. Furthermore, Yeo *et al.*'s [14] meta-analysis focused solely on all-cause dementia. Our study incorporated the most recent literature and conducted a meta-analysis of multiple types of dementia, building on the findings of these three previous meta-analyses. This approach allowed us to more thoroughly assess the relationship between cataract surgery and dementia risk. Compared to earlier studies, our research is innovative in several ways. First, our meta-analysis included the latest literature and increased the sample size, enhancing the reliability and credibility of the findings. Second, we analyzed different types of dementia, rather than aggregating the results of different types. Finally, to explore the potential influencing factors, we conducted subgroup analysis and sensitivity analysis, which were not addressed in previous meta-analyses.

Results from a meta-analysis of five cohort studies included in this research indicated that cataract surgery was not associated with a re-

duced risk of all-cause dementia in cataract patients. After excluding the study by Lee *et al.* [6], it was found that cataract surgery was indeed associated with a reduced risk of all-cause dementia in cataract patients. This was consistent with the results of previous meta-analyses [12–14]. We hypothesize that this discrepancy may be due to the inclusion of other eye diseases, such as glaucoma, diabetic retinopathy, and age-related macular degeneration, in the multifactorial analysis, as these conditions are known risk factors for dementia and cognitive impairment. Overall, cataract surgery is associated with a lower risk of dementia in cataract patients. The APOE $\epsilon 4$ mutation is the most important genetic risk factor for Alzheimer's disease [15–17]. In this meta-analysis, a subgroup analysis was performed to determine whether APOE $\epsilon 4$ mutations were adjusted. We found that after adjusting for APOE $\epsilon 4$ mutations in two studies [7, 8], cataract surgery was associated with a significantly reduced risk of all-cause dementia in cataract patients. Meanwhile, a meta-analysis that included the same two studies [7, 8] found that cataract surgery was associated with a reduced risk of Alzheimer's disease in cataract patients. Therefore, this study strongly supports the conclusion that cataract surgery can help reduce the risk of Alzheimer's disease in cataract patients.

Cataract is a treatable visual impairment of the eye [18]. Visual impairment may increase the risk of dementia and cognitive impairment [3, 19, 20]. Early prevention and timely treatment of visual impairment can reduce the incidence of dementia [21]. Surgery can significantly reduce the risk of all-cause dementia and Alzheimer's disease in cataract patients. Researchers have analyzed the mechanisms of cataract surgery and reduced dementia risk. First, after cataract surgery, people can receive higher-quality sensory input, which may help mitigate the risk of dementia [6]. Second, cataract surgery reactivates cells that are sensitive to blue light. A specific group of cells in the retina, known as intrinsically photosensitive retinal ganglion cells (ipRGC), are particularly sensitive to blue light stimulation and play a crucial role in regulating circadian rhythms [22, 23]. Degeneration and functional alterations in these cells have been shown to be associated with cognition and Alzheimer's disease [23–25]. Because cataract surgery restores the passage of blue light through the lens into the ipRGC, it reactivates these cells to prevent cognitive decline [7].

Limitations of this study. First, it was a cohort-based analysis, and the adjusted covariates in the results varied, which may introduce potential confounding factors. Consequently, the influence of other variables on dementia risk cannot be entirely ruled out. Second, the study was based

on observational data, and randomized controlled trials are needed to validate the association between cataract surgery and dementia risk. Third, due to the limited number of included studies, there might be potential publication bias. Fourth, the included studies only compared cataract patients who underwent surgery with those who did not. Fifth, while the number of participants in the study was substantial, it may not be representative of the broader population. Participants may be from specific regions or specific medical institutions, which could limit the generalizability of the findings. Lastly, cognitive impairment and dementia are complex conditions influenced by various factors, including genetics, vascular disease, and inflammation. Therefore, it is unclear whether cataract surgery directly causes dementia or if other mediating variables are involved. As a result, the question of whether cataracts represent an independent risk factor for dementia requires further investigation.

In conclusions, the available evidence suggests that cataract surgery may reduce the risk of Alzheimer's disease in cataract patients. However, the relationship between cataract surgery and the reduced risk of all-cause dementia in these patients requires further investigation. The study had several limitations, including its observational design, potential confounding factors, limitations in sample characteristics, and inability to determine causation. Future studies will need to employ more rigorous study designs and explore specific mechanisms to further validate this association and provide more robust evidence.

Funding

No external funding.

Ethical approval

Not applicable.

Conflict of interest

The authors declare no conflict of interest.

References

- Salardini A. An overview of primary dementias as clinicopathological entities. *Semin Neurol* 2019; 39: 153-66.
- Xiong Z, Li X, Yang D, Xiong C, Xu Q, Zhou Q. The association between cataract and incidence of cognitive impairment in older adults: a systematic review and meta-analysis. *Behav Brain Res* 2023; 450: 114455.
- Jefferis JM, Mosimann UP, Clarke MP. Cataract and cognitive impairment: a review of the literature. *Br J Ophthalmol* 2011; 95: 17-23.
- Tamura H, Tsukamoto H, Mukai S, et al. Improvement in cognitive impairment after cataract surgery in elderly patients. *J Cataract Refract Surg* 2004; 30: 598-602.
- Yoshida Y, Ono K, Sekimoto S, Umeya R, Hiratsuka Y. Impact of cataract surgery on cognitive impairment in older people. *Acta Ophthalmol* 2024; 102: e602-11.
- Lee C, Shin E, Kim M, et al. The effect of cataract surgery on the risk of dementia: a nationwide cohort study. *J Clin Med* 2023; 12: 6441.
- Lee CS, Gibbons LE, Lee AY, et al. Association between cataract extraction and development of dementia. *JAMA Inter Med* 2022; 182: 134-41.
- Ma LZ, Zhang YR, Li YZ, et al. Cataract, cataract surgery, and risk of incident dementia: a prospective cohort study of 300,823 participants. *Biol Psychiatry* 2023; 93: 810-9.
- Miyata K, Yoshikawa T, Morikawa M, et al. Effect of cataract surgery on cognitive function in elderly: results of Fujiwara-kyo Eye Study. *PLoS One* 2018; 13: e0192677.
- Yu WK, Chen YT, Wang SJ, Kuo SC, Shia BC, Liu CJ. Cataract surgery is associated with a reduced risk of dementia: a nationwide population-based cohort study. *Eur J Neurol* 2015; 22: 1370-7, e1379-80.
- Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010; 25: 603-5.
- Liu X, Guan Z, Liang S, Feng S, Zhou Y. Associations of cataract, cataract surgery with dementia risk: a systematic review and meta-analysis of 448,140 participants. *Eur J Clin Invest* 2024; 54: e14113.
- Zhang Q, Ju Y, Zheng W, et al. Association of cataract extraction and the risk of dementia-a systematic review and meta-analysis. *Front Aging Neurosci* 2023; 15: 1168449.
- Yeo BSY, Ong RYX, Ganasekar P, Tan BKJ, Seow DCC, Tsai ASH. Cataract surgery and cognitive benefits in the older person: a systematic review and meta-analysis. *Ophthalmology* 2024; 131: 975-84.
- Williams-Gray CH, Goris A, Saiki M, et al. Apolipoprotein E genotype as a risk factor for susceptibility to and dementia in Parkinson's disease. *J Neurol* 2009; 256: 493-8.
- Elias-Sonnenschein LS, Viechtbauer W, Ramakers IH, Verhey FR, Visser PJ. Predictive value of APOE-ε4 allele for progression from MCI to AD-type dementia: a meta-analysis. *J Neurol Neurosurg Psychiatry* 2011; 82: 1149-56.
- Talyansky S, Le Guen Y, Kasireddy N, Belloy ME, Greicius MD. APOE-ε4 and BIN1 increase risk of Alzheimer's disease pathology but not specifically of Lewy body pathology. *Acta Neuropathol Commun* 2023; 11: 149.
- Song P, Wang H, Theodoratou E, Chan KY, Rudan I. The national and subnational prevalence of cataract and cataract blindness in China: a systematic review and meta-analysis. *J Glob Health* 2018; 8: 010804.
- Cao GY, Chen ZS, Yao SS, et al. The association between vision impairment and cognitive outcomes in older adults: a systematic review and meta-analysis. *Aging Mental Health* 2023; 27: 350-6. 3
- Pellegrini M, Bernabei F, Schiavi C, Giannaccare G. Impact of cataract surgery on depression and cognitive function: systematic review and meta-analysis. *Clin Exp Ophthalmol* 2020; 48: 593-601.
- Shang X, Zhu Z, Wang W, Ha J, He M. The association between vision impairment and incidence of dementia and cognitive impairment: a systematic review and meta-analysis. *Ophthalmology* 2021; 128: 1135-49.
- Kwok CPC, Kwok JOT, Yan RWK, et al. Dementia and risk of visual impairment in Chinese older adults. *Sci Rep* 2022; 12: 18033.

23. Xiao Z, Wu W, Zhao Q, Liang X, Luo J, Ding D. Association of glaucoma and cataract with incident dementia: a 5-year follow-up in the shanghai aging study. *J Alzheimer's Dis* 2020; 76: 529-37.
24. Von dem Hagen EA, Houston GC, Hoffmann MB, Jeffery G, Morland AB. Retinal abnormalities in human albinism translate into a reduction of grey matter in the occipital cortex. *Eur J Neurosci* 2005; 22: 2475-80.
25. Vandewalle G, Maquet P, Dijk DJ. Light as a modulator of cognitive brain function. *Trends Cognitive Sci* 2009; 13: 429-38.