

Diagnostic value of methylated SEPT9 and RNF180 for gastric cancer in high-risk population: a multi-centre retrospective study in Yunnan, Southwest China

Ying Huang¹, Shu Yang², Yao Cun³, Juying He⁴, Guihua Xu⁵, Pinghui Shi⁶, Jinmei Yang^{7*}, Qiang Wang^{8*}

¹Mengzi People's Hospital, Mengzi, China

²Southern Central Hospital of Yunnan Province, Gejiu, China

³The First People's Hospital of Xundian Hui and Yi Autonomous County, Kunming, China

⁴Chenggong District People's Hospital, Kunming, China

⁵Kaiyuan People's Hospital, Kaiyuan, China

⁶Gejiu People's Hospital, Gejiu, China

⁷First Affiliated Hospital of Kunming Medical University, Kunming, China

⁸The Affiliated Hospital of Yunnan University, Kunming, China

***Corresponding authors:**

Jinmei Yang
First Affiliated
Hospital of Kunming
Medical University
Kunming, China

Qiang Wang
The Affiliated Hospital
of Yunnan University,
Kunming, China

Submitted: 11 October 2024; **Accepted:** 3 December 2024

Online publication: 31 December 2024

Arch Med Sci

DOI: <https://doi.org/10.5114/aoms/196807>

Copyright © 2024 Termedia & Banach

High mortality from gastric cancer (GC) is a serious challenge to global health, especially in East Asia and Eastern Europe, where GC remains one of the leading causes of cancer-related deaths [1]. Early detection of GC tends to improve treatment efficiency and survival rates. Currently available early detection methods mainly involve endoscopic biopsy, which is considered the gold standard of diagnosis. Other than these, noninvasive techniques include a serum pepsinogen test and a *Helicobacter pylori* (Hp) test as a screening technique for the high-risk population. Recently, ways have critically emphasised some molecular markers based on methylated DNA with specific microRNAs presented in blood and gastric juice, which are expected to give better rates of early detection [2]. Given their detectability in gastric mucosal carcinogenesis, methylated SEPT9 and ring finger protein 180 (RNF180) are acquiring prominence in the field of GC early diagnosis. The SEPT9 gene is already known to deal with the cytoskeletal tissue, and its methylation was considered to have a bright future in the blood-based detection methodology that shall not be invasive for the screening of GC [3]. Also, the RNF180 gene, encoding an E3 ubiquitin-protein ligase, also shows methylation changes in gastric tissue detection in tissue samples and blood [4]. The detection of combining these 2 biomarkers is thought to increase the accuracy of diagnosis, providing options for the early detection and management of GC [5]. Therefore, we seek to determine their effects in diagnosing GC patients in a Chinese population and compare diagnostic performance across all gastric diseases and cancer stages.

This was a retrospective study. The study involved the retrieval of stored blood samples and medical documentation from individuals who had undergone endoscopic biopsy for early GC screening between January 2021 and December 2023. High-risk subjects were defined as aged ≥ 40 years, resident in regions where the incidence of stomach cancer is high. Those with Hp and those who had a previous history of pre-cancerous gastric conditions like atrophic chronic gastritis, gastric ulcer, gastric

polyp, or pernicious anaemia were included. This study followed the Standards for Reporting of Diagnostic Accuracy (STARD) reporting guideline.

The clinical data obtained encompassed demographic details such as age and gender, as well as comprehensive clinical characteristics including the type and location of gastric lesions, Lauren classification, TNM staging, and lymph node metastasis status. Additionally, we quantified key tumour markers using the Roche E-70 analyser. To differentiate between benign and malignant gastric conditions, specific cut-off values were established: ≥ 5 ng/ml for CEA, ≥ 37 U/ml for CA19-9, and ≥ 6.9 U/ml for CA72-4.

Because this *SEPT9/RNF180* methylation detection process uses a fixed approach, processes were conducted under fixed circumstances to control and maintain the validity and reliability of the results. In line with this, 10 ml blood samples were drawn up in K2EDTA anticoagulant tubes from a BD Vacutainer® to avoid clotting and centrifuged for 12 min under 1350 ± 150 rpm so that

the plasma could separate. If not to be immediately processed, the plasma was stored between 2°C and 8°C. In detail, plasma was centrifuged for the second time under the same conditions, 3.5 ml aliquots were placed in labelled conical tubes, and then they were stored at -25 to -15°C for DNA preservation for a period of 2 weeks.

DNA methylation was assessed using a two-step process employing the PCR fluorescence probe technique. Initially, free DNA was isolated from plasma and treated with bisulfite to convert unmethylated cytosines to uracil sulfonate, leaving methylated cytosines unaltered. Subsequently, triplex PCR amplification was conducted, in which specific PCR blockers and fluorescent probes identified and amplified methylated sequences of the *RNF180* and *SEPT9* genes. Each sample, along with positive and negative controls, was processed in parallel to ensure consistency.

These statistical analyses were performed with Python 3.12. The receiver operating characteristic (ROC) curves and the area under the ROC curve (AUC) were calculated to assess the performance of the different biomarkers as a diagnostic test. Estimation of confidence intervals for sensitivity, specificity, positive predictive value, and negative predictive value was done via the efficient score method as provided in statsmodels Wilson score interval.

In this study, 8 hospitals in Yunnan Province were selected. The blood samples of 201 patients who underwent endoscopic biopsy for early GC screening were retrospectively collected from these hospitals from January 2021 and December 2023: 78 with GC (GC+) and 123 without GC (GC-). The GC+ group had a mean age of 60.2 years (SD ± 5.3), whereas the GC- group had a mean age of 57.6 years (SD ± 5.9). The baseline characteristics are summarised in Table I.

Figure 1 A revealed that methylated *SEPT9* and *RNF180* had varying positive detection rates across cancer stages, with 43.2% in early stages and 58.8% in late stages. In non-cancerous cases, these markers demonstrated a positive detection rate of 13.0%. Figures 1 B and 1 C present ROC curves quantifying the opportunity of *SEPT9/RNF180* detection by different GC groups and markers. Figure 1 D displays the ROC curves for detection using different combinations of biomarkers. Various characteristics of GC were assessed for its favourable detection rates, presenting a significant difference by the type and location of GC (Figure 1 E). The ROC curve categorised by age and gender highlights differences in diagnostic performance (Figure 1 F).

The findings from this study underscore the significant potential of methylated *SEPT9* and *RNF180* as biomarkers for the detection of gastric cancer. The study demonstrates that both

Table I. General characteristics of study population

Variable	GC+	GC-
Participant, <i>n</i>	78	123
Age [years] mean \pm SD	60.2 \pm 5.3	57.6 \pm 5.9
Gender, <i>n</i> (%)		
1 Male	47 (60.3)	64 (52.0)
0 Female	31 (39.7)	59 (48.0)
Gastric lesions, <i>n</i> (%)		
Atrophic gastritis	–	48 (39.0)
Intestinal metaplasia	–	19 (15.4)
Gastric polyps	–	22 (17.9)
Gastric ulcer	–	34 (27.7)
Lauren classification, <i>n</i> (%)		
Intestinal type	41 (52.6)	–
Diffuse type	25 (32.1)	–
Other	12 (15.3)	–
Tumour location, <i>n</i> (%)		
Body	29 (37.2)	–
Fundus	32 (41.0)	–
Pylorus	17 (21.8)	–
TNM staging, <i>n</i> (%)		
I	18 (23.1)	–
II	26 (33.3)	–
III	32 (41.0)	–
IV	2 (2.6)	–
Lymph node metastasis, <i>n</i> (%)		
Yes	11 (14.1)	–
No	67 (85.9)	–

GC+ – gastric cancer patients; GC- – non-gastric cancer patients.




Figure 1. Diagnostic value of methylated SEPT9 and RNF180 for gastric cancer. **A** – Positive detection rates of methylated SEPT9 and RNF180 in participants at different stages. **B** – ROC curves for SEPT9/RNF180 detection. **C** – ROC curves for various gastric cancer markers. **D** – ROC curve comparison of combined biomarkers. **E** – Positive detection rates for gastric cancer characteristics. **F** – ROC curves by age and gender

biomarkers exhibit high specificity and improved detection rates in later stages of the disease, with the positive detection rate escalating from 43.2% in early stages to 58.8% in advanced stages of gastric cancer. The detection results of dif-

ferent biomarker combinations revealed that the combination of SPET9 and RNF180 achieved the highest diagnostic accuracy. Notably, the intestinal type of gastric cancer demonstrated the highest detection rate at 75.6%, indicating that

these biomarkers are particularly effective in identifying more aggressive forms of the disease. Furthermore, the study highlights a clear distinction in diagnostic performance based on patient demographics, with males and older individuals showing significantly higher detection accuracies, which could inform tailored screening approaches in clinical settings.

Utilising peripheral blood for cell-free DNA (cfDNA) methylation testing has emerged as a pivotal non-invasive diagnostic method for early cancer detection. Studies have shown that methylation variations in colorectal cancer tissues may indicate the potential effectiveness of *SEPT9* methylation in peripheral blood for early CRC screening [6]. This came about with the ‘Shield’ early screening tool that applies changes in cfDNA genomic variations and DNA methylation status and fragmentation patterns for colorectal cancer. The recently completed ECLIPSE study reported sensitivity and specificity of diagnosing CRC at 83.1% and 89.6%, respectively [7].

Further studies have revealed that high *RNF180* and *SEPT9* methylation in the plasma samples of patients with gastric cancer were sound and reasonable biomarkers for that neoplasm [8]. *RNF180* is a tumour suppressor gene that suppresses gastric cancer cells’ growth, proliferation, and migration. Methylation in its promoter region, however, is positively associated with cancer cell growth, differentiation, and metastasis. *RNF180* is one of the hypermethylated genes detected in the plasma of people with gastric cancer. Similarly, the *SEPT9* gene, another tumour suppressor, regulates apoptosis and cell division. The sensitivity and specificity of *SEPT9* methylation for diagnosing gastric cancer were 47.7% and 92.3%, respectively. However, for complex gastric lesions, these findings fell short of the required diagnostic accuracy. Combining methylated *SEPT9* with protein markers could significantly improve diagnostic sensitivity [9].

Our results agree with the fact that DNA methylation markers, being ideal gastric adenocarcinoma biomarkers, are more sensitive than traditional protein tumour markers such as CEA, CA199, and CA125 for gastric cancer detection [10]. Notably, studies have demonstrated that DNA methylated biomarkers such as *SEPT9* and *RNF180* are particularly effective in distinguishing between malignant and non-malignant gastric tissues with high specificity. This supports our findings, which indicate specificity rates exceeding 85% for both biomarkers [11]. In addition, the differential performance of each biomarker across various cancer stages and types in our study concurs with Xu *et al.*, who found enhanced detection in more advanced stages of gastric cancer [12]. Our work

is new to the literature as we quantitatively compared the different detection capabilities according to patients’ demographics, showing significant variations likely to make a big difference in screening strategy.

Nevertheless, several limitations deserve consideration. The small sample size, drawn from only 8 hospitals in a single geographic region (Yunnan, Southwest China) limits the generalisability of the findings to broader populations. In contrast, blood-based biomarkers, which are less invasive to patients compared to tissue biopsies, may carry far less assessment in the whole heterogeneity of gastric cancer compared to tissue research, which may affect the sensitivity and specificity in different subtypes and at various stages of the tumour. Furthermore, it failed at this stage to engage in a longitudinal follow-up of this biomarker appraising its prognostic value.

In conclusion, we found that methylated *SEPT9* and *RNF180* are effective biomarkers for the detection of GC, especially in advanced stages and specific GC types. Their incorporation into clinical practice has the potential to improve early detection and patient prognosis in high-risk populations. However, this conclusion requires further confirmation from clinical studies with larger samples.

Acknowledgments

Thank you to Ms. Zhao Xiuyan (Baoshan People’s Hospital) for her contribution to the manuscript revision.

Ying Huang and Shu Yang contributed equally to this work.

Funding

No external funding.

Ethical approval

This study was approved by the Ethics Committee of the First Affiliated Hospital of Kunming Medical University (No. FEY-ZD-24-2.19).

Conflict of interest

The authors declare no conflict of interest.

Reference

1. Smyth EC, Nilsson M, Grabsch HI, van Grieken NC, Lordick F. Gastric cancer. *Lancet* 2020; 396: 635-48.
2. Roy S, Kanda M, Nomura S, et al. Diagnostic efficacy of circular RNAs as noninvasive, liquid biopsy biomarkers for early detection of gastric cancer. *Mol Cancer* 2022; 21: 42.
3. Zhao L, Li M, Zhang S, Liu Y. Plasma-methylated *SEPT9* for the noninvasive diagnosis of gastric cancer. *J Clin Med* 2022; 11: 6399.

4. Shen SY, Singhania R, Fehringer G, et al. Sensitive tumour detection and classification using plasma cell-free DNA methylomes. *Nature* 2018; 563: 579-83.
5. Wang X, Dong Y, Zhang H, et al. DNA methylation drives a new path in gastric cancer early detection: current impact and prospects. *Genes Dis* 2023; 11: 847-60.
6. Lofton-Day C, Model F, Devos T, et al. DNA methylation biomarkers for blood-based colorectal cancer screening. *Clin Chem* 2008; 54: 414-23.
7. Chung DC, Gray DM 2nd, Singh H, et al. A cell-free DNA blood-based test for colorectal cancer screening. *N Engl J Med* 2024; 390: 973-83.
8. Zhang Q, Du Z, Wang X, et al. Cell-free nucleic acid as promising diagnostic biomarkers for gastric cancer: a systematic review. *J Cancer* 2024; 15: 2900-12.
9. Tang H, Ma X, Zhou L, et al. Distinct performance of methylated SEPT9 in upper and lower gastrointestinal cancers and combined detection with protein markers. *Genet Test Mol Biomarkers* 2022; 26: 239-48.
10. Nie Y, Gao X, Cai X, et al. Combining methylated SEPTIN9 and RNF180 plasma markers for diagnosis and early detection of gastric cancer. *Cancer Commun (Lond)* 2023; 43: 1275-9.
11. Cao CQ, Chang L, Wu Q. Circulating methylated Septin 9 and ring finger protein 180 for noninvasive diagnosis of early gastric cancer. *Transl Cancer Res* 2020; 9: 7012-21.
12. Xu J, Song J, Wang T, et al. A combination of methylation and protein markers is capable of detecting gastric cancer detection by combined markers. *Epigenomics* 2021; 13: 1557-70.